

1 DELIVERY SYSTEMS FOR MYCOTECHNOLOGIES, MYCOFILTRATION
2 AND MYCOREMEDIATION

3 This application is a continuation-in-part of U.S. patent application
4 serial no. 09/790,033 for DELIVERY SYSTEMS FOR
5 MYCOTECHNOLOGIES, MYCOFILTRATION AND MYCOREMEDIATION,
6 filed 2/20/2001, currently co-pending, herein incorporated in its entirety by
7 reference.

8 BACKGROUND OF THE INVENTION

9 1. Field of the Invention

10 The present invention is generally related to products and methods for
11 inoculation with beneficial fungi. More particularly, the present invention is
12 related to the use of fungal slurries, landscaping cloths, paper products and
13 mats, hydroseeding equipment and agricultural equipment for inoculation
14 with spores and hyphae of mushrooms and other fungi for purposes including
15 ecological rehabilitation and restoration, bioremediation, habitat preservation
16 and agriculture.

17 2. Description of the Related Art

18 The foundation and continuation of life is directly dependent upon
19 healthy habitats. Habitats are increasingly in peril due to the expansion of
20 human enterprises, exacerbating the effects of erosion, and leading to losses
21 in biodiversity and ecological resilience. In the construction of roads,

1 expansion of suburbia and urban centers, trees and shrubs are removed and
2 topsoils are stripped away and soils are compacted. As rains ensue, the forces
3 of erosion further threaten ecological health in removing latent soils and
4 causing sediment accumulation in the lowlands. This severe loss of topsoil
5 tenacity directly results in enormous expenses both societally and
6 environmentally. Certain human enterprises have also resulted in the
7 contamination of widespread areas with toxic wastes and pollutants.

8 The vegetative, long-lived body of a fungus is an extensive network of
9 microscopic threads (known as mycelium, mycelia or mycelial hyphae) which
10 fully permeates soil, logs, or others substrates within which the organism
11 grows. Most ecologists now recognize that soil health is directly related to the
12 presence, abundance and variety of fungal associations. The mycelial
13 component of topsoil within a typical Douglas fir forest in the Pacific
14 Northwest approaches 10% of the total biomass; the threadlike hyphae of
15 fungal mycelia may exceed one mile of mycelium per cubic inch of soil.
16 Healthy ecosystems include a wide variety of fungal associations. For
17 example, mycorrhizal fungi (including many mushroom fungi) form a
18 mutually dependent, beneficial relationship with the roots of host plants,
19 ranging from trees to grasses to agricultural crops. When the mycelia of
20 these fungi form an exterior sheath covering the roots of the plant they are
21 termed ectomycorrhizal; when they invade the interior root cells of host

1 plants they are called endomycorrhizal (also known as vesicular-arbuscular
2 or VA mycorrhizae). Saprophytic fungi (wood and organic matter
3 decomposers) are the primary decomposers in nature, working in concert with
4 a succession of microorganisms and plants to break down and recycle organic
5 and inorganic compounds and materials. Saprophytic fungi have also been
6 found to form symbiotic, mutually beneficial relationship with a number of
7 agricultural crops. For example, corn is known to give bigger yields in the
8 presence of straw bales inoculated with *Stropharia rugosoannulata* as
9 compared to uninoculated straw bales. The no-till method of farming also
10 benefits from the growth of Basidiomycetes including mushrooms, reducing
11 plant stubble into nutrients. Parasitic mushrooms have their own role in a
12 healthy ecosystem, although they can become overly destructive in unhealthy
13 systems. Another broad class of decomposers is the more primitive, non-
14 mushroom forming "fungi imperfecti," including also molds and yeasts.

15 Evidence of the premier role of fungi as decomposers can easily be
16 gathered in a walk through a healthy forest--rotting logs that have been
17 infested by fungi. Without the presence of fungi, few if any organisms are
18 able to effectively degrade the complex aromatic polymers cellulose and
19 lignin, the two primary components of woody plants; cellulose, and
20 particularly lignin, the most recalcitrant of substrates in nature, are
21 generally otherwise resistant to microbial attack and decomposition. The

1 fungi, particularly "white rot fungi," which are adept at decomposing lignin,
2 and "brown rot fungi," premier decomposers of cellulose, produce a complex
3 suite of enzymes that oxidize the structures completely to water and carbon
4 dioxide via a radical-mediated mechanism.

5 Both liquid substrate and solid substrate cultures of white rot fungi
6 have been the subject of years of bioremediation research in numerous
7 laboratories, as evidenced by the large number of publications and patents in
8 this area. See, for example, U.S. Patent Nos. 4,554,075 (1985), 4,891,320
9 (1990), 5,085,998 (1992), 5,486,474 (1996), 5,583,041 (1996) and 5,597,730
10 (1997). Such saprophytic white rot wood-decomposing fungi have shown the
11 ability to degrade recalcitrant foreign compounds such as polynuclear
12 aromatic hydrocarbons (PAHs), alkanes, creosote, pentachlorophenol (PCP),
13 polychlorinated biphenyls (PCBs), dichlorodiphenyltrichloroethane (DDT),
14 trinitrotoluene (TNT), dioxin, nitrogenous compounds such as ammonium
15 nitrate, urea, purines and putriscines, as well as agricultural wastes and
16 agricultural runoff. However, these bioremediation processes have
17 significant limitations, hindering the transition from the laboratory to large
18 scale field applications, and in general have not been used commercially. One
19 particular problem has been that economic and effective delivery systems for
20 large scale field applications of white rot fungi have not been available.

21 The saprophytic fungi have also proven to be efficient digesters of

1 potentially harmful organisms such as coliform bacteria and nematodes. The
2 voracious Oyster mushrooms (*Pleurotus ostreatus*) have been found to be
3 parasitic against nematodes. Extracellular enzymes act like an anesthetic
4 and stun the nematodes, thus allowing the invasion of the mycelium directly
5 into their immobilized bodies.

6 For these and other reasons there has been great interest in fungi for
7 uses such as introduction of mycorrhizal fungi, bioaugmentation of soils,
8 bioremediation, biological control and production of mushrooms.

9 Among the methods for delivering fungal spores and hyphal inoculum
10 to soil for various purposes such as bioremediation or agriculture are carriers
11 such as grain, sawdust and wood chip spawn, alginate hydrogels with and
12 without additional nutrient sources, vermiculite and peat optionally
13 saturated with nutrient broths, vermiculite and rice flour or grain flour,
14 straw or other agricultural waste products overgrown with fungal mycelium,
15 pelleted fungal inoculum preparations, etc.

16 The usual methods for inoculation with fungi are typically expensive,
17 labor intensive and/or ineffective. Various techniques have been used to
18 inoculate growing substrates with those fungi known as mushrooms. These
19 include methods of inoculating beds of wood chips, beds of compost, lawns and
20 soils. Also known are methods of inoculating soils with fungi for the
21 bioremediation purposes.

1 Beds of wood chips are typically inoculated by spreading sawdust
2 and/or woodchip spawn (spawn being defined herein as any material
3 inoculated with mycelium or impregnated with mycelium and used for
4 inoculation) throughout the wood chips or by placing a layer of spawn within
5 the wood chips. Beds of compost are typically inoculated in a similar manner
6 with a grain spawn, although a sawdust spawn may also be utilized in some
7 instances. The use of expensive spawn of limited shelf life produced by labor-
8 and equipment-intensive sterile culture methods are among the
9 disadvantages of this approach.

10 Another method of inoculation involves spore mass inoculation or
11 inoculation with mycelia fermented under sterile conditions. In the first
12 method spores may be collected and broadcast, but more preferably is
13 conducted by immersion of the mushroom(s) in water to create a spore mass
14 slurry, the addition of molasses, sugars and/or sawdust to stimulate spore
15 germination, aeration, incubation and broadcast of the aqueous spore mass
16 slurries. This approach and the similar approach with liquid mycelium
17 inoculated and grown under sterile conditions may be successfully utilized.
18 These approaches, however, require either fresh spore-producing mushrooms
19 or sterile culture techniques, and application must be during the time frame
20 of vigorous peak growth after germination or inoculation or the mycelial
21 fragments will not coalesce into a contiguous mycelial mat. There remains a

1 need for more convenient products and processes for widespread application
2 of biologically active spore and/or mycelial inocula.

3 Trees, lawns and seedbeds have been inoculated with mycorrhizal
4 species using various tablets or gels prepared from spores or mycelium. Trees
5 may also be inoculated with mycorrhizal mushrooms by dusting the roots of
6 seedlings with spores or mushroom mycelium or by dipping the exposed roots
7 of seedlings into water enriched with the spore mass of the mycorrhizal
8 species. Another method for inoculating mycorrhizae calls for the planting of
9 young seedlings near the root zones of proven mushroom-producing trees,
10 allowing the seedlings to become 'infected' with the mycorrhizae of a
11 neighboring tree. After a few years, the new trees are dug up and
12 transplanted. Another method involves broadcasting spore mass onto the
13 root zones of trees. Such approaches can be labor intensive, expensive, of
14 uncertain success and/or not suited to widespread use.

15 Patented approaches for inoculation with mycorrhizal fungi include
16 U.S. Patent No. 4,294,037 (1981) to Mosse *et al.* for a process for the
17 production of vesicular-arbuscular (VA) mycorrhizal fungi comprising
18 growing a VA fungus on plant roots in nutrient film culture for 1 to 3 months
19 and harvesting for inoculum production; U.S. Patent No. 5,178,642 (1993) to
20 Janerette for culturing of ectomycorrhizal fungal inoculants on a solid
21 medium, contacting the mycelia in the solid medium with perlite wetted with

1 a nutrient solution, incubating for about three months and broadcasting; and
2 U.S. Patent No. 4,551,165 (1985) to Warner for mycorrhizal seed pellets
3 formed from vesicular-arbuscular mycorrhizal inoculum peat, at least one
4 seed and a binder compacted into pellet form. It is also known to add various
5 compositions to seeds to assist growth. For example, U.S. Patent No.
6 5,586,411 (1996) to Gleddie *et al.* discloses methods for adding *Penicillium*
7 *bilaii* and *Rhizobium* bacteria in a sterilized peat base to legume seeds so as
8 to increase the availability of soluble phosphate and fixed nitrogen. However,
9 it is not known to add mycorrhizal fungi directly to seeds, nor is it known to
10 combine saprophytic or entomopathogenic fungi directly with seeds or
11 seedlings, nor is it known to combine mycorrhizal fungi with saprophytic,
12 entomopathogenic and/or imperfect fungi for the purpose of habitat
13 restoration. Again, there remains a need for cheaper and more efficacious
14 methods for large scale use.

15 U.S. Patent No. 6,033,559 discloses microbial mats constructed of
16 stratified layers of cyanobacteria and purple autotrophic bacteria, and
17 optionally other microorganisms such as algae or fungi, organized into a
18 layered structure held together with slime with an organic nutrient source
19 provided, optionally with support structures such as shredded coconut hulls,
20 ground corn cobs or wood fiber. While such bacterial mats may be suited to
21 aquatic environments, they are not particularly suited for terrestrial

1 applications. An additional disadvantage is that algae are generally not as
2 'enzymatically equipped' to deal with toxins and pollutants, the fungi being
3 the keystone species which render nutrients available to the photosynthetic,
4 chlorophyll producing algae and plants.

5 Trends in spawn technology have long been evolving towards pelletized
6 or granular spawn, for purposes such as inoculation of substrates for
7 production of gourmet and medicinal mushrooms, inoculation with
8 mycorrhizal fungi, inoculation with white rot fungi for bioremediation and
9 inoculation with fungi imperfecti for control of soilborne pathogens. Various
10 forms of pelletized spawn are known, including those formed from nutrients,
11 with or without binders, and peat moss, vermiculite, alginate gel, alginate gel
12 with wheat bran and calcium salts, hydrophilic materials such as hydrogel,
13 perlite, diatomaceous earth, mineral wool, clay, etc. See Stamets, *Growing*
14 *Gourmet and Medicinal Mushrooms* (1993) and U.S. Patent Nos. 4,551,165
15 (1985), 4,668,512 (1987), 4,724,147 (1988), 4,818,530 (1989), 5,068,105
16 (1991), 5,786,188 (1998) and 6,143,549 (2000). Pelletized spawn is
17 specifically designed to accelerate the colonization process subsequent to
18 inoculation. Examples of pelletized spawn range from a form resembling
19 rabbit food to pumice-like particles.

20 Idealized pelletized spawn seeks a balance between surface area,
21 nutritional content, and gas exchange and enables easy dispersal of mycelium

1 throughout the substrate, quick recovery from the concussion of inoculation,
2 and sustained growth of mycelium sufficient to fully colonize the substrate.
3 Many grains and other substrates are, however, pound-for-pound, particle for
4 particle, more nutritious than most forms of pelletized spawn. Furthermore,
5 use of grains or liquid-inoculum or other forms of inoculum avoids the
6 expense and labor of pelletizing. There remains a need for more economical
7 and more efficacious means of inoculation of large scale areas.

8 It is known that berms and revetments and other protective structures
9 are employed to halt soil erosion caused by runoff or precipitation. One
10 particular, well-known system for the creation of such protective structures
11 consists in the construction and use of "gabions," *e.g.*, "mattress gabions,"
12 large, thin rectangular containers filled with gravel, crushed stone and other
13 material, fitted with a cover and consisting of galvanized or galvanized and
14 plastic-coated wire netting panels joined together with ties or wire stitches
15 and designed to cover, without any break, extensive tracts of land of the most
16 disparate conformation, as if they were actual 'mattresses.' Similar
17 structures may be constructed of "basket gabions," "sack gabions," "gabion
18 mats" and "log gabions."

19 In many applications, there is a need for gabions to rehabilitate the
20 environment and allow development of an ecosystem able to utilize the water
21 runoff, thereby resisting erosion in a more environmentally sound manner.

1 In other applications, a gabion that is biodegradable would be more useful
2 than those metal or other degradation-resistant materials used to construct
3 gabions. There is also a need for gabions that could 'filter' contaminants such
4 as agricultural runoff, including fertilizer, animal waste and pesticide runoff,
5 urban runoff, etc. for protection of streams and rivers. In many situations
6 there is also a need for gabions of cheaper materials.

7 There is, therefore, a continuing need for enhancing the effectiveness of
8 fungal inoculation and growth and thereby improving habitat preservation
9 and habitat recovery. There is also a need for enhanced products and
10 methods for accomplishing fungal inoculation as an aid to such and habitat
11 recovery and preservation. There is also a need for such fungal products and
12 methods as an aid to agriculture, including both plant cultivation and
13 mushroom cultivation.

14 In view of the foregoing disadvantages inherent in the known types of
15 fungal inoculants, the present invention provides improved inoculating
16 agents and methods of using such agents.

17 BRIEF SUMMARY OF THE INVENTION

18 Fungi have been found by the present inventor to be a "keystone
19 species," one that facilitates a cascade of other biological processes that
20 contribute to healthy ecologies, the fungi being necessary for health of
21 environments and capable of "leading the way" in the remediation,

1 reclamation, restoration and/or preservation of environments. As fungi,
2 including many or all gourmet and medicinal saprophytic mushroom fungi,
3 produce extracellular enzymes and acids not only capable of breaking down
4 cellulose and lignin, but also hydrocarbons such as oils, petroleum products,
5 fuels, propellants, PCBs and many other pollutants, the fungi are
6 particularly suited to bioremediation of badly polluted and eroded
7 environments, depleted environments, etc. Such fungi have also been found
8 to be a keystone in the most healthy and luxuriant terrestrial environments.
9 Fungal organisms are now known as the largest biological entities on the
10 planet, with various individual mats covering more than 20,000 acres,
11 weighing 10,000 kg. (22,000 lb.) and remaining genetically stable for more
12 than 1,500 years. The momentum of mycelial mass from a single mushroom
13 species, growing outwards at one-quarter to two inches per day, staggers the
14 imagination. These silent mycelial tsunamis affect all biological systems
15 upon which they are dependent. As one fungus matures and dies back, a
16 panoply of other fungi come into play, acting to catalyze habitat recovery and
17 habitat health.

18 Nearly all plants have joined with saprophytic and mycorrhizal fungi
19 in symbiosis. Plants may devote a majority of the net energy fixed as
20 sunlight to below ground processes, not only root growth but also to feed
21 mycorrhizal fungi and other microorganisms. However, this symbiotic

1 relationship is not a net energy loss. Mycorrhizal fungi surround and
2 penetrate the roots of grasses, shrubs, trees, crops and other plants,
3 expanding the absorption zone ten- to a hundred-fold, aiding in plants' quest
4 for water, transferring and cycling macro and micro nutrients, increasing soil
5 aeration and the moisture-holding capacity of soils and forestalling blights,
6 pathogens and disease. With the loss of fungi, the diversity of insects, birds,
7 flowering plants and mammals begins to suffer, humidity drops, now-exposed
8 soils are blown away, and deserts encroach. To aid in the solution of these
9 problems, new "mycotechnologies" (after mycology, the study of fungi) are
10 provided herein.

11 In view of the disadvantages inherent in the known products and
12 methods for fungal inoculation, the present invention provides improved
13 products and methods for intensive and/or widespread inoculation of
14 beneficial fungal species. The present invention provides new products and
15 methods utilizing fungal spore and hyphal compositions, useful for
16 impregnation of soils, fabric landscaping cloths, soil blankets and rugs, mats,
17 mattings, bags, gabions, fiber logs, fiber ropes, fiber bricks, etc.; useful for
18 distribution via spray hydroseeding equipment and mobile hydroseeders;
19 useful for agricultural planting equipment, harvesting equipment and field
20 preparation equipment; useful for cultivation of gourmet and medicinal
21 mushrooms; and useful for the habitat restoration and preservation uses

1 described herein. Inoculation with beneficial fungal spores and/or mycelial
2 hyphae, and optionally and preferably with seeds, provides products and
3 methods useful for purposes including enhancing plant growth and
4 mycorrhizal and symbiotic relationships, habitat restoration, erosion control
5 and stabilization of soils, treatment of contaminated habitats, filtration
6 (“mycofiltration”) of agricultural and urban water runoff, fungal
7 bioremediation (“mycoremediation”) of biological and chemical pollutants and
8 toxic wastes, and production of mycelia and mushrooms and improved
9 production of plants, providing nutrients to insects, herbivores and numerous
10 organisms up and down the food chain. Preferred fungi include the “fungi
11 perfecti” (including those fungi producing gilled and polypore and other
12 mushrooms) and the “fungi imperfecti” (the simpler, non-mushroom
13 producing fungi including molds and yeasts) and their various forms of
14 mycelium and spores, including both sexually produced and asexually
15 produced spores and spore variations. Particularly useful are the saprophytic
16 mushrooms for purposes such as mycoremediation and mycofiltration of
17 agricultural and urban runoff, the saprophytic and mycorrhizal fungi for
18 improvements in agricultural products and methods, the entomopathogenic
19 fungi for insect control, and combinations of the saprophytic, mycorrhizal,
20 entomopathogenic and/or other fungi imperfecti. Such products and methods
21 further provide reduced costs, ease of application and improved efficiency

1 when compared to known products and processes.

2 The fungal inoculation products and the fungal methods of the present
3 invention may, depending upon the application, advantageously include
4 habitat recovery and restoration, erosion control, rapid decay and
5 decomposition of forest debris and agricultural waste, bioremediation of
6 contaminated sites through decomposition of hydrocarbon based
7 contaminants and concentration/removal of heavy metals from soils,
8 adjustment of soil pH, mycofiltration of agricultural and industrial runoff,
9 large-scale introduction of mycorrhizal species, gourmet species and other
10 beneficial mushroom species, introduction of entomopathogenic (capable of
11 causing disease in insects) fungi for control of pest insects, fungi for control of
12 soilborne plant pathogens, the production of gourmet and medicinal
13 mushrooms, and numerous other applications. A water-spore, water-
14 mycelial hyphae or water-spore and/or hyphae-seed slurry (or similar slurries
15 with vegetable or other oils) may be applied directly to soils. Alternatively,
16 the water-spore, water-mycelial hyphae or water-spore-hyphae or oils
17 suspension is applied to commercially available products such as landscaping
18 cloths, gabions, mats, burlap and other fiber bags, paper and/or cardboard
19 materials, bulk substrates or other fiber substrates, etc., optionally
20 simultaneously with or followed by seed application. As another alternative,
21 such products may be inoculated by traditional inoculation methods, such as

1 those utilizing grain spawn or sawdust spawn. Less preferably, similar
2 products made of non-biodegradable materials may be utilized. A water-seed-
3 spore mass or water-seed-mycelial hyphae slurry offers a novel approach for
4 inoculating environments with fungi and can be applied directly to bare soils,
5 straw, reeds, wood chips, sawdust, fibers and fiber products, landscape
6 fabrics and papers, burlap sacks, gabions, etc. The mycelial hyphae may be
7 utilized fresh, dried or freeze-dried. The benefits of these products and
8 approaches include ease of application, erosion control, habitat restoration,
9 mycofiltration, mycoremediation, and mycorrhizal and fungal associations.

10 The use of such aforementioned fungally impregnated biodegradable
11 membranes, in combination with plant seeds allows for a unique delivery
12 system: cardboard boxes whose side walls have been infused or applied with
13 plant seeds in combination with fungal spores, mycelium, or extracts of the
14 mycelium of mycorrhizal, symbiotic, saprophytic, and entomopathogenic
15 fungi. A multiplicity of problems are solved with one solution. The
16 prevalence of cardboard boxes delivered throughout the world on a daily basis
17 exceeds thousands of tons per day, boggling the imagination. The cardboard
18 box is ubiquitous to the world community. The predominance of cardboard in
19 the manufacturing of boxes and its over-abundance strains the resources of
20 communities. With this invention, cardboard boxes have a value-added, after
21 market benefit as they become a living resource for ecological recovery. The

1 panels of the box can be used for home gardening, commercial agriculture, for
2 mycofiltration, mycoremediation, and mycopesticidal purposes. The box can
3 be used as an educational tool for teaching children while at the same time be
4 the container for transporting items related or unrelated to the invention.
5 The cardboard boxes become an ecological footprint for creating a garden,
6 seed bed, an orchard, a forest and even an expanding oasis, starting the
7 process of habitat improvement and recovery. An added advantage is that
8 the cardboard panels can be placed over soil to suppress competitive weed
9 growth and to retain moisture. The decomposition of the paper based
10 materials by the fungus releases nutrients to aid plant growth.

11 Oils may also be used as a carrier material. Petroleum oils can be
12 readily digested by certain fungi and biodegradable oils are readily digested
13 by most or all fungi perfecti and fungi imperfecti. Therefore oil-spore or oil-
14 hyphae mixtures or water-oil-spore or water-oil-hyphae suspensions, with or
15 without seeds, provide an alternative to the water-spore or water-hyphae
16 slurries which may be utilized in the practice of the present invention. In
17 general, biodegradable oils are preferred as offering an environmentally
18 friendly and a more readily available nutritional source to a wide variety of
19 fungi. Such fungal or hyphal oils may also be preferably employed in
20 applications such as ecological rehabilitation, mycoremediation and
21 mushroom growing where use of a vegetable oil as an additional nutritional

1 source is desired.

2 The use of fungi as keystone organisms releases nutrients into the
3 surrounding environment from the biodegradable carrier materials to
4 enhance the growth of targeted or naturally occurring plants, from grasses to
5 shrubs to trees to complex biological communities. In essence, biological
6 successionism can be directed through the use of a single species or a complex
7 plurality of fungal components, using fungi as the keystone organisms
8 leading the way in habitat enhancement or recovery. The fungi may
9 optionally be used in combination with plants, algae, lichen, bacteria, etc.

10 Biodegradable fabric cloths and blankets made of straw, coconut fibers,
11 corn stalks, wood fibers and other similar materials, wood chips and straw
12 bales are in common use along roadsides to help prevent or lessen erosion
13 and help ecological recovery. When plant root growth increases in these
14 locations, the tenacity of the soil is enhanced, lessening the chances for
15 erosion. However, none use a fungal component as a determining factor in
16 enhancing the effects of such biodegradable erosion-control materials. The
17 present invention offers improved products wherein fungi act as a “keystone”
18 or “linchpin” species, ameliorating the impact of erosive forces by helping to
19 establish communities of organisms, using fungi to enhance or control the
20 growth of other organisms including but not limited to plants, protozoa,
21 bacteria, viruses, algae, lichens, invertebrates, arthropods, worms and/or

1 insects. Also advantageous is the use of fungal mycelium to enhance the
2 tenacity of overlaying fabric cloths or bulk substrate on habitats, thus
3 preventing 'slippage' and anchoring the fabric cloths, wood chips, straw, etc.

4 Such mycelial products are also useful for combating viruses and
5 virulent bacteria, for example *Escheria coli*, *Bacillus subtilis*, malaria,
6 cholera, anthrax, and water-borne diseases, as well as biological warfare
7 (BW) pathogenic species. By infusing mycelium into cloths, blankets,
8 gabions, mats, berms, etc., targeted disease organisms such as bacteria,
9 fungi, viruses, protozoa and amoebas can be effectively reduced, ameliorating
10 the downstream impact as well as in residence. Such benefits could help
11 fisheries, for instance, stave off *Pfiesteria*.

12 In another embodiment of the present invention, fungal spores and/or
13 mycelial hyphae are introduced into hydroseeding equipment, agricultural
14 seeding equipment, harvesting equipment and other agricultural equipment.
15 This allows for the simultaneous inoculation of beneficial fungi directly into
16 lawns, disturbed soils, agricultural fields, agricultural wastes, etc.

17 The addition of fungal tissue (spore mass and/or hyphae) into
18 landscaping materials, hydroseeding-type equipment and all types of
19 agricultural equipment is an effective means for the simultaneous replanting
20 and fungal inoculation of disturbed or recovering environments, leading to
21 habitat restoration, improved control of runoff and mycofiltration of runoff

(trapping biological and chemical contaminants, denaturing them), etc. The addition of fungal inocula to agricultural equipment can provide improved means of introducing beneficial symbiotic saprophytic fungi and mycorrhizal fungi, entomopathogenic fungi for control of insect pests and fungi imperfecti for control of soilborne plant pathogens. Introduction of such fungal inocula into harvesting equipment can provide efficient means of inoculating agricultural waste products or efficient production of inoculated straw bales and rounds, etc., useful for the practice of many embodiments of the present inventions.

Another advantage of the present invention lies in the use of fungal components to accelerate the decomposition of biodegradable fabrics and other materials in sensitive environments where such fabrics and materials have been placed for the purposes of preventing erosion and enhancing habitat recovery.

Another advantage of the present invention arises from the use of fungal components in biodegradable materials to enhance water retention properties of such materials, using the natural water-absorption properties of mycelium.

Supplementary advantages arise from the fact that fungally colonized mycelial fiber substrates liberate carbon dioxide, essential for healthy plant growth, especially essential for young seedlings. As the grass or other plants

1 grow up, it creates a high humidity layer through condensation formation
2 from dew point as well as the 'greening' effect which is naturally cooler.

3 Further advantages arise from the use of adsorbent or absorbent
4 biodegradable fiber cloths and mats inoculated with spores and/or hyphae of
5 petroleum oil-eating fungi. Thus the oil slicks or spills may be soaked up by
6 the cloth or mat material and digested by the mycelium of the fungus.

7 An additional advantage is the use of fungally impregnated
8 biodegradable materials along stream and sensitive watersheds to ameliorate
9 the impact of runoff containing sediment and pollutants. The use of such
10 products allows for sequestration of excess or harmful nitrogenous,
11 phosphorus-laden or carbonaceous compounds as well as sediment and silt
12 from gravel roads and other sources. Fisheries, especially spawning streams
13 of salmon and trout, as well as other species such as shellfish, benefit directly
14 and dramatically from mycofiltration of silt and sediment, which can create
15 an environment inhospitable to eggs, and pollutants, which can have far-
16 ranging negative effects. Numerous advantages naturally follow the use of
17 such mycelial products and methods to protect sensitive watersheds such as
18 salmon and trout spawning grounds, riparian runoff and wetlands, thereby
19 providing mushroom and mycelial biomass which then feeds developing
20 larvae of numerous insects, providing additional benefit to fisheries and
21 recreational users through enhancement of the food chain as well as through

1 protection from upland runoff.

2 The present invention provides further advantages via use of a fungal
3 component or components in biodegradable materials to help catalyze
4 significant climate change in arid environments through the enhancement of
5 the water retention capacities of the top soils, leading to the 'oasis'
6 phenomena in dryland habitats, the net effects of which are not only erosion
7 control, but significant enhancement of biological communities which then
8 can become 'seed' banks leading to a creations of satellite communities in
9 proximity to the genome source.

10 Another advantage of the present invention is the use of fungal
11 components in biodegradable materials to create communities of fungi,
12 including commercially valuable mushrooms.

13 Additional advantages arise when such products and methods are used
14 to bioremediate contaminated, toxic and hazardous sites, providing
15 breakdown of dangerous organic, inorganic and biological threats while
16 simultaneously triggering the ecosystem recovery as above. In biologically
17 hostile environments, a small sample of the targeted habitat can be
18 introduced to the fermentation of the fungal mycelia, at a late stage, so that
19 the chosen fungal candidate can acclimate to the complex biota of the
20 targeted environment. This technique reduces transplant shock, and further
21 enhances the effectiveness of the present invention.

Further advantages arise from the use of colonized fiber substrates to combat virulent bacteria, reduce or eliminate viruses, limit pathogenic fungi, yeasts, and molds, control protozoa such as amoebas, ciliates, flagellates, and sporozoans, control multicellular organisms such as rotifers and trap and digest nematodes.

Further advantages are obtained when such 'mycocloths' and 'mycomats' are infused with fungi capable of decomposing biological and chemical warfare toxins. The mycocloths and mycomats can then be used to decontaminate toxic landscapes, battlefield and otherwise, thus leading to reuse of valuable land.

Still further advantage may be gained from use of fungally impregnated biodegradable materials, either contained within or in the absence of a matrix of biodegradable or non-biodegradable materials, to concentrate heavy metals, for example radioactive metals and precious metals, which then can be removed to eliminate toxins topically and subsurface. Such residual organic debris and mycelia could be economically or profitably separated from the metals through incineration, biodigestion with other organisms (*e.g.*, bacteria, protozoa or yeasts) and or via chemical treatments (*e.g.*, enzymes, acids or catalysts).

The present invention provides further advantages through use of entomopathogenic fungal components to control, reduce or eliminate pest

1 insects or disease-carrying insects in the applied environments. Extracts of
2 the pre-conidial mycelium of entomopathogenic fungi may also be utilized to
3 attract and/or control insects. More broadly, fungal components in
4 biodegradable materials may be utilized to control harmful insects, enhance
5 insect communities, or invite beneficial insects in the applied environments.
6 Since insect communities can influence or predetermine bird communities,
7 the fungal constituent has a direct downstream effect on this and many other
8 biological successions.

9 The present invention thus allows for wide scale inoculation of desired
10 mushroom species on widely varying substrates suitable for use in various
11 applications and environments. Numerous advantages arise from growing
12 beneficial fungi and mushrooms for various agricultural, forestry, ecological
13 and bioremediation purposes including habitat restoration and preservation,
14 rapid decay of forestry byproducts and wastes, mycofiltration of agricultural
15 and industrial runoff, decomposition of hydrocarbon based contaminants and
16 toxins, concentration/removal of heavy metals from soils, sewage or other
17 substrates, insect, pest and disease control, soil improvement and adjustment
18 of soil pH, introduction of mycorrhizal fungi, production of gourmet and
19 medicinal mushrooms, improved crop yields, etc.

20 The present invention has been found to achieve these advantages.
21 Still further objects and advantages of this invention will become more

1 apparent from the following detailed description and appended claims.
2 Before explaining the disclosed embodiments of the present invention in
3 detail, it is to be understood that the invention is not limited in its application
4 to the details of the particular products and methods illustrated, since the
5 invention is capable of other embodiments which will be readily apparent to
6 those skilled in the art. Also, the terminology used herein is for the purpose
7 of description and not of limitation.

8 DETAILED DESCRIPTION OF THE INVENTION

9 Innovations of the present invention include introducing saprophytic
10 fungi, mycorrhizal fungi, entomopathogenic fungi, fungi imperfecti and/or
11 other fungi as keystone species using a wide variety of novel products and
12 methods. By infusing substrates or soils with fungal inoculum as disclosed
13 herein, widespread areas of land, sensitive areas such as stream banks and
14 riparian areas, drainages into wetlands, areas in need of topsoil
15 supplementation, polluted areas, etc. may be favorably treated and
16 transformed via fungi. By selecting the type of fungal spores or hyphae to be
17 infused, an ecologist, remediator, forester, farmer, landscaper and others can
18 direct the course of ecological recovery or ecological preservation, thereby
19 improving the economical usefulness of the land for varying forest, farm,
20 riparian, agricultural and urban uses. Furthermore, by selecting the types of
21 seeds, persons can further direct the course of development--for example, by

1 using a mixture of grasses and trees, the grasses typically germinating first
2 followed by germination of the tree seeds. Alternatively, seedlings may be
3 directly utilized. Such fungal inoculation may be accomplished via fibrous
4 fabrics, hydroseeding equipment or a variety of agricultural equipment.

5 In one embodiment, spores, spore mass, actively growing mycelial
6 hyphae, dried or freeze dried powdered fungal mycelium, and/or powdered
7 mushroom fruitbodies are placed into carrier materials used for landscaping
8 and ecological purposes. Mycorrhizal fungi and/or various wood, lawn and
9 field mushrooms and/or entomopathogenic fungi and/or fungi imperfecti may
10 be utilized. The landscaping carrier materials are preferably also
11 impregnated with the seeds of grasses, native grasses, flowers, native
12 wildflowers, and/or trees and other plants. Although some seeds may become
13 'fungi food,' particularly when fresh live mycelium is utilized, some seeds will
14 survive and germinate. Alternatively, such landscaping carrier products may
15 be inoculated, overgrown with mycelium, and seeds then added. Additional
16 organisms such as bacteria, lichens, moss, algae, etc., as well as other fungi,
17 both perfect and imperfect, may optionally be added. Such mats or larger
18 fabrics or other fiber products may be overlaid onto disturbed grounds both to
19 aid plant growth and as a vehicle for treating contaminated habitats, wherein
20 the mycelium acts as a mycofiltration membrane, trapping biological and
21 chemical contaminants and denaturing them. Similarly, a wide variety of

1 landscaping carrier products, discussed in more detail below, may similarly
2 be utilized. The present invention also includes kits for the construction of
3 such fabrics, mats and other fiber carrier products.

4 Mycomaterials which are utilized after being overgrown with mycelium
5 may be utilized fresh or metabolically arrested via refrigeration for storage
6 and transport. Alternatively, the mycelium may be metabolically arrested
7 through freeze-drying (flash chilling), drying, or by other means, for storage,
8 transportation and subsequent rehydration for field deployment. Storage
9 time of up to a year or more is possible. It will be understood that such
10 metabolic arresting of development may encompass either a slowing of
11 metabolism and development (such as refrigeration) or a total suspension or
12 shutdown of metabolism (freeze-drying, air-drying and cryogenic suspension).

13 The novel fungal inoculum/seed sprays and slurries may be applied
14 directly to soils. For many applications it is preferable to apply fungal
15 inoculum to landscaping materials such as wood or straw bulk substrates,
16 mulches, biodegradable landscaping fabrics and blankets, mats, bags,
17 gabions, fiber baskets, fiber-logs, fiber-bricks, cardboard, paper, etc., thereby
18 providing an initial nutritional source, particularly in applications such as
19 habitat restoration, erosion control, mycoremediation, mycofiltration,
20 landscaping, etc.

21 The mycotechnologies of the present invention may be utilized in the

1 various states of fungal lifecycle, with or without seeds. Where a landscaping
2 type application is desired, a preferred embodiment will often be a paper,
3 cardboard or fabric cloth-seed-spore and/or mycelial hyphae embodiment,
4 with germination of spores, hyphae and seeds occurring upon placement and
5 watering or rainfall. Such may also be preferred in certain erosion control
6 and habitat preservation or rehabilitation applications. For other
7 applications, such as mycoremediation, berm building and mushroom
8 cultivation, mycocloths overgrown with live fungal mycelium on thicker, more
9 rug-like or mat-like materials may sometimes be preferred. For these and
10 other applications, it may be preferable to form a fibrous material, such as
11 burlap, into a sack or bag, or to form a thicker material into bags, basket
12 gabions or mattress gabions and fill with woody fiber and/or non-woody fiber
13 materials. Such sacks, bags and gabions, and optionally their contents, may
14 be inoculated with spores, fresh mycelial hyphal fragments, dried or freeze-
15 dried mycelial hyphae, powdered mushrooms or spawn or combinations
16 thereof, and utilized either immediately after inoculation or after the fibrous
17 material has been overgrown by hyphae, depending on circumstances and
18 desired use. The mats may be deployed in various settings, including both
19 terrestrial and aquatic (such as floating mats). Mycomaterials which are not
20 initially combined with seeds may later have seeds or growing plants added,
21 for combined efficacy with the fungal component for bioremediation, erosion

1 control, landscaping aesthetics, etc.

2 Suitable landscaping and/or non-landscaping materials, carriers and

3 spawn products include geocloths and geofabrics, soil blankets, landscaping

4 fabrics and other fabrics, nettings, rugs, mats, mattings, fiber felt pads, straw

5 tatamis, mattress inserts, burlap bags, papers, fiber logs, fiber bricks,

6 gabions, cardboards, papers, etc. These materials, carriers and products may

7 be formulated of any suitable fiber, including those derived from woody and

8 non-woody fibers such as wood chips, sawdust, wood pulp, wood mulch, wood

9 wastes, leaf paper, wood-based papers, non-wood papers, pressed cardboard,

10 corrugated cardboard, fiberized rag stock, cellophane, hemp and hemp-like

11 materials, bamboo, papyrus, jute, flax, sisal, coconut fibers, wheat straw, rice

12 straw, rye straw, oat straw and other cereal straws, reeds, rye grass and

13 other grasses, grain hulls and other seed hulls such as cottonseed hulls,

14 cornstalks, corncobs, soybean roughage, coffee plant waste and pulp, sugar

15 cane bagasse, banana fronds, palm leaves, the hulls of nuts such as almonds,

16 walnuts, sunflower, pecans, peanuts, etc., soy waste, cactus waste, tea leaves

17 and the wide variety of other agricultural waste products and combinations

18 thereof. Suitable animal fibers include wool, hair and hide (leather) and

19 combinations thereof. In general, biodegradable wood or plant fibers are

20 preferred over non-biodegradable synthetic fibers. Such is particularly the

21 case with fabrics, mats, blankets, bags, gabions, fiber-logs, etc. utilized for

1 purposes such as mycoremediation, mycofiltration, construction of
2 biodegradable berms, levees, revetments, embankments, etc. Suitable
3 synthetic fibers include plastics and polymers such as polypropylene,
4 polyethylene, nylon, etc. The fibrous woody and non-woody plant fibers may
5 be in any form including paper, textile, fabric, veil, mat, matted, mesh
6 matting, matting rug, felt pressing, blanket, filter, woven, woven roving, open
7 weave, nonwoven, knitted, strand roving, continuous strand, chopped strand,
8 knotted, yarn, braided ropes, milled fiber, high-pressure extrusion rope or
9 mat, composites, etc. and combinations thereof.

10 Carrier materials may optionally be amended to provide additional
11 nutrients via spraying or soaking of the materials in sugars such as maltose,
12 glucose, fructose or sucrose, molasses, sorghum, mannitol, sorbitol, corn steep
13 liquor, corn meal and soybean meal, vegetable oils, casein hydrolysate, grain
14 brans, grape pumice, ammonium salts, amino acids, yeast extract, vitamins,
15 etc. and combinations thereof. Typically such amendments should be utilized
16 sparingly or with materials that are to be pasteurized or sterilized, as such
17 amendments, particularly carbohydrates and nitrogen supplements, may
18 greatly reduce substrate semi-selectivity for fungi and increase the risk of
19 contamination after fungal inoculation.

20 Carrier materials such as cardboard panels or other paper-based
21 membranes, can be inoculated with fungi and plant seeds. Such panels can

1 be incorporated into the manufacturing of boxes, especially cardboard boxes.
2 If mycorrhizal, saprophytic and/or mycopesticidal fungi are used in concert
3 with compatible seeds of plants, the cardboard panels become springboards
4 for life and ecological recovery. Fibers selecting from the group consisting of
5 paper pulp fibers, cellophanes (including those with silicon fibers), shredded
6 paper products, wood fibers, sawdusts, corn, jute, coir, coconut, hemp, wheat,
7 rice, grasses, coffee, cotton, kenaf, mosses, lichens, mugworts, wools, animal
8 skins, and biodegradable polymers can also be utilized for the construction of
9 membranes or box panels incorporating this invention. The aforementioned
10 materials can be reformulated to incorporate fungi in the form of spores or
11 mycelium in combination with plant seeds. The boxes still serve their
12 traditional, structural function for the delivery of goods, but now have
13 increased value for their after-delivery use. The panels or boxes could be used
14 for other purposes unrelated to this invention, and increased value because of
15 its further utility in growing plants, enhancing food production and for
16 bioremediation. The panels of the box host assortments of seeds customized
17 to the ecological and cultural specifics of their destination. The selection of
18 seeds predetermines the selection of mycorrhizal and saprophytic fungi. Upon
19 unpacking the box's contents, the box is disassembled by hand or by sharp
20 instrument. The cardboard panels, infused with seeds and fungi, are laid
21 upon or into soil. With the addition of water, the cardboard softens, the fungi

1 are activated, and the seeds germinate. Immediately upon germination the
2 seeds have contact with beneficial fungi, insuring an early symbiotic
3 relationship before competitor fungi can harm the seeds. The mycorrhizal
4 fungi stimulate shoot and root growth, expand the sphere of the root zone for
5 absorption of water and nutrients, improve the micro-hydrology of the
6 surrounding soil, and protect the young plants from diseases. With moisture,
7 the saprophytic fungi decompose the cardboard, freeing more nutrients. The
8 cardboard layer lessens evaporation, preserves moisture, shades and cools the
9 soil underneath. The softening cardboard allows the penetration of the shoots
10 and roots. If the cardboard is scored with fine cuts during manufacturing, the
11 roots and shoots can emerge unencumbered. The cardboard fully
12 decomposes, becoming soil, and leaves no waste.

13 One of the many useful applications of this 'living box', that is, a box
14 constructed with dormant fungi and seeds, for assisting refugees, indigenous
15 displaced peoples, including victims from natural and man-made disasters.
16 As the first emergency relief often is delivered to refugees in a box, there is
17 the economically feasible opportunity of utilizing the delivery box as inoculum
18 for growing plants and fungi. The insides of the box could be sorted according
19 to species of plants, climatic zones, pH requirements, and soil conditions. By
20 example but not by limitation, the seeds of the plant species could be selected
21 from the group comprising of corn, wheat, rice, oats, rye, lentils, beans,

1 squash, melons, potatoes, carrots, turnips, garlic, ginger, mustard, chard,
2 cilantro, fennel, oregano, chives, basil, thyme, and onions. Such box panels
3 would be recognized by the recipients as having a value, a natural currency
4 for anyone who has an interest in cultivating and habitat recovery. The
5 educational lesson from having children using the 'living box' is as important
6 an advantage of this invention as any aspect previously described.

7 The use of cloths, rugs, mats, papers, cardboards, etc. for fungal
8 inoculation products and methods makes advantageous use of several fungal
9 characteristics. For example, it has been found by the present inventor that
10 quite different techniques are called for when inoculating soils and non-
11 sterile substrates as compared to sterile substrates. When inoculating
12 sterilized or pasteurized substrates, or materials composted so as to prepare a
13 selective nutritious medium of such characteristics that the growth of
14 mushroom mycelium is promoted to the practical exclusion of competitor
15 organisms (see *The Mushroom Cultivator* (1983) by Stamets and Chilton), a
16 technique known as "through spawning" is preferable, wherein the fungal
17 inoculum is introduced via numerous inoculation points (such as colonized
18 grain spawn or sawdust spawn) throughout the medium. However, such an
19 approach in non-sterile bulk substrates such as wood chips or soil may lead to
20 disaster. Each inoculation point becomes a separate colony surrounded by
21 competitor organisms in all directions, often with the result that the

1 inoculation points are unable to generate the necessary mycelial momentum
2 to successfully colonize the substrate. The present inventor has found “layer
3 spawning” or “sheet inoculation,” wherein the fungal inoculum is spread in a
4 horizontal layer within the non-sterile bulk substrate, to be much more
5 successful. Such sheet inoculation takes advantage of several fungal
6 characteristics: 1) mycelia often grows and spreads most rapidly in the
7 lateral, horizontal directions; 2) when mycelia grows horizontally and links
8 into a mycelial layer or mat, it becomes much more vigorous, resistant to
9 contaminants and competitive, allowing further successful growth and
10 colonization in the vertical direction; and 3) ‘wild’ mycelial organisms are
11 typically matlike and layered in that they may cover many acres, yet be only
12 a few inches deep. Thus a landscaping cloth or mat introduces inoculation
13 points and allows for horizontal growth in accord with the mushroom or
14 fungi’s natural characteristics. By having a contiguous sheet of mycelium
15 above toxins, extracellular enzymes can “rain” down, effectively decomposing
16 them.

17 It has further been found that when “sandwich inoculation” utilizing
18 two or more such layers of inoculum is utilized, competitiveness and ultimate
19 success is even further enhanced as the two mycelial layers grow vertically
20 and link up, forming a thoroughly colonized block. In such cases, having two
21 (or more) layers of fungal inoculum with substrates sandwiched in between

1 gives more resilience, allowing for more duration, increasing effectiveness
2 over the long term. Thus when mycelial landscaping cloths or mycelial mats
3 are preferred, a plurality of mats or cloths in stacked, separated layers will
4 often be even more preferable. It will be noted that when cloths are formed
5 into a bag or sack, inoculated with spores or hyphae, and filled with bulk
6 substrates such as woodchips, two lateral layers of cloth are naturally
7 formed, plus a route for initial vertical growth and linkup of layers is
8 provided. Thus in many application, such 'mycobags' will be preferred. Such
9 mycobags and similar mattress gabions, preferably filled with wood chips,
10 straw, composts, agricultural waste products, etc., are also particularly useful
11 for building biodegradable erosion control structures, berms, revetments,
12 banks, barriers, dykes, retaining wall structures, channel liners, filter drain
13 systems, etc. for purposes such as mycofiltration and mycoremediation. It
14 will also be noted that heavy cloths may be formed into 'basket gabions'
15 which will also provide multiple horizontal layers for growth and routes for
16 vertical colonization when stacked to form revetments, berms, barriers,
17 banks, etc. In general, biodegradable cloths are preferred, but non-
18 biodegradable materials such as plastic polymers may also be inoculated and
19 utilized as an inoculation source for non-sterile bulk substrates. Such
20 mycomats, mycocloths, mycobags and mycogabions may be treated with
21 fungal inocula for immediate use or may be partially overgrown or completely

1 overgrown with fungi and then utilized. In many cases, seeds are also
2 preferably added, such as native grasses, etc. The use of burlap (typically
3 made of jute, flax or hemp) mycobags filled with wood chips on 'mineral
4 earth,' the layer beneath topsoil, has also been found to be an effective way to
5 begin the process of soil regeneration.

6 The use of cardboard, straw, sawdust, etc. layers on top of the
7 inoculated materials (such as bags, blankets, cloths, etc.) or substrate
8 material is useful to ameliorate the loss of water, whether these inoculated
9 materials are overlaid on the ground or buried under wood chips, straw or
10 agricultural waste products. For example, layers of cardboard (top), wood
11 chips (middle), and inoculated cloth or bag (bottom), or alternatively
12 cardboard (top), inoculated cloth (middle) and wood chips (bottom) or
13 variations thereof. The use of moisture retaining materials on top is also
14 useful when 'sandwich' layers of inoculated materials and uninoculated
15 substrate are utilized. Ultimately, the insulating material itself will be
16 transformed in a rich soil.

17 In order to increase fungal penetration of soils, berms, etc. beyond the
18 typical 10-20 cm. (4-8 inch) depth, aeration methods or oxygenated water
19 may be employed. Various methods of aeration and oxygenating water and
20 delivering such will be readily apparent to those skilled in the art. By way of
21 example but not of limitation, water may be oxygenated by means of

1 percolation, high pressure infusion, electrolysis, hydrogen peroxide, chemical
2 reaction, etc.

3 Where it is desired to use fungally inoculated and enhanced
4 landscaping cloths, mats, gabions, fiber-logs, fiber-bricks or bulk substrates of
5 a size or amount that exceeds even the size of the largest autoclaves (for
6 pressure steam sterilization) or steam pasteurization chambers, or where
7 steam sterilization or pasteurization is not available, the various alternative
8 methods known to the art may be utilized. By way of example but not of
9 limitation, these methods include: 1) Immersion of the landscaping cloth or
10 other substrate in a hydrated lime (calcium hydroxide) solution, thereby
11 largely rendering competitor fungi and bacteria inactive from the drastic
12 change in pH. For example, 2-4 pounds of lime is added for every 50 gallons
13 of water, resulting in a lime/water ration of about .5%-1.0%. The cloth or
14 substrate is soaked overnight or for a similar period, the water is drained and
15 the cloth or substrate is inoculated using standard spawn methods or
16 methods as disclosed herein. Such is particularly useful for fungi that can
17 tolerate an alkaline environment better than competitors, such as *Pleurotus*.
18 Optimizing the parameters for the species being cultivated, such as initial pH
19 of the makeup water, greatly influences the success or failure of this method.;
20 2) Immersion of the cloth or substrate in a bleach bath utilizing household
21 bleach (typically about 5.25% sodium hypochlorite). For example, 3-4 cups of

1 household bleach is added for every 50 gallons of water, the cloth or bulk
2 substrate is immersed and kept submerged for a minimum of 4 and a
3 maximum of 12 hours, and the bleach leachate is drained off. The cloth or
4 bulk substrate is immediately inoculated.; or 3) Disinfection with hydrogen
5 peroxide (H_2O_2). This technique has been refined by Rush Wayne, who,
6 having become frustrated with the difficulty and expense of creating a sterile
7 environment in his home, refined this technique to a practical level. A full
8 description of this technique can be found at
9 *www.members.aol.com/PeroxyMan* and detailed instructions may be found in
10 the book *Growing Mushrooms the Easy Way: Home Mushroom Cultivation*
11 *with Hydrogen Peroxide* by R. Wayne (1999), Rush Wayne Enterprises,
12 Eugene, Oregon, herein incorporated by reference. It should be noted,
13 however, that much resident contamination can survive this process. While
14 hydrogen peroxide works to kill many fungal spores, yeasts and bacteria by
15 producing a reactive form of oxygen, which destroys cell walls, because fungal
16 compounds have evolved to decompose organic compounds in the
17 environment using peroxides and peroxidases, the mycelia of contaminant
18 fungi and molds is protected from its oxidizing effects. If colonies of mycelium
19 from contaminant fungi have already developed, this method will be of
20 limited advantage. Although not thorough enough to neutralize most of the
21 natural fungi contaminants resident in raw sawdust, straw, etc., hydrogen

1 peroxide can help complete the process started with many preheated
2 substrates. For example, when wood is baked in an oven at 149°C (300°F) for
3 3 hours, compounds are destroyed in the wood that would otherwise
4 neutralize the peroxide. Hydrogen peroxide can be diluted 100-fold, from 3%
5 to .03%, into water (less than 60°C or 140°F). This water can then drench the
6 substrate to further reduce the likelihood of competitors.; 4) High-pressure
7 extrusion of straw and sawdust and other bulk substrates. This method for
8 treating straw and sawdust utilizes the heat generated from the extrusion of
9 a substrate from a large orifice through a smaller one, producing pellets or a
10 'rope' substrate. The effective reduction of the substrate causes frictional
11 heat to escalate. For example, a 6:1 reduction of straw into a 10 millimeter
12 pellet creates a thermal impaction zone where temperatures exceed 80°C
13 (176°F), temperatures sufficient for pasteurization. Alternatively, a roller
14 mechanism may be utilized rather than a narrow orifice, enabling processing
15 of much more substrate mass and producing a matlike product.; 5) The
16 detergent bath method, which utilizes biodegradable detergents containing
17 fatty oils to treat bulk substrates. Coupled with surfactants that allow
18 thorough penetration, these detergents kill a majority of the contaminants
19 competitive to mushroom mycelium. The landscaping cloth, mat or bulk
20 substrate is submerged into and washed with a detergent solution. The
21 environmentally benign wastewater is discarded, leaving the cloth, mat or

1 substrate ready for inoculation.; and 6) A yeast fermentation method may be
2 utilized to render straw and other substrates suitable. Straw can be
3 biologically treated using yeast cultures, specifically strains of beer yeast,
4 *Saccharomyces cerevisiae*. This method by itself is typically not as effective
5 as those previously described. First, a strain of beer yeast is propagated in
6 200 liters (~50 gallons) warm water to which malt sugar has been added (for
7 example, 1-5% sugar broth). Fermentation proceeds for 2 to 3 days
8 undisturbed in a sealed container at room temperature. Another yeast
9 culture can be introduced for secondary, booster fermentation that lasts for
10 another 24 hours. After this period of fermentation, chopped straw or other
11 substrate is forcibly submerged into the yeast broth for no more than 48
12 hours. Not only do these yeasts multiply, absorbing readily available
13 nutrients, which can then be consumed by the mushroom mycelium, but
14 metabolites such as alcohol and antibacterial byproducts are generated in the
15 process, killing competitors. Alternatively, the natural resident microflora
16 from the bulk substrate may be utilized for submerged fermentation. After 3
17 or 4 days of room-temperature fermentation, a microbial soup of great
18 biological complexity evolves. The broth, which can be used as a natural
19 biocide, is now removed and the substrate is inoculated. Although highly
20 odiferous for the first 2 days, the offensive smell soon disappears and is
21 replaced by the sweet fragrance of actively growing mycelium. The outcome

1 of any of these alternative methods greatly depends on the cleanliness of the
2 substrate being used, the water quality, the spawn rate, and the aerobic state
3 of the medium during colonization. These alternative methods generally do
4 not result in the high consistency of success (>95%) typical with heat
5 treatment techniques.

6 It will be noted that normally paper rolls, paper towels, cardboard, etc.
7 are 'clean' enough and structurally selectively favors the fungal mycelium so
8 that products constructed of such may be utilized without pasteurization or
9 sterilization (especially cardboard such as corrugated or pressed cardboard).

10 Where prior sterilization of the ground is desired, the many various
11 methods known to the art may be utilized, for example flame, hydrogen
12 peroxide, hydrogen peroxide/acetic acid, etc.

13 In another preferred embodiment, fungal inoculum is added to spray
14 hydroseeding equipment or mobile landscaping hydroseeders for delivery of
15 spores and/or hyphae.

16 Where non-pasteurized or non-sterilized large fabrics or geocloths,
17 including wire mesh reinforced erosion control cloths and synthetic fabrics,
18 are, for example, used for landscaping, used to stabilize soil embankments,
19 slopes and walls, used to promote vegetation growth while providing rockfall
20 protection and/or used for mycofiltration or mycoremediation, a preferred
21 embodiment is 'spray hydroseeding' of fungally inoculated products. Spray

1 hydroseeding is performed with a pump for dense liquids, which sprays on to
2 the surface to be greened a mixture consisting of, for example, fungal inocula
3 (spores, dried hyphae, powdered mushrooms, conidia, etc.), seeds, fertilizer if
4 desired, and commercial green hydromulch (a wood fiber mulch) or soil
5 improvement substances, optionally and usually preferably with a binder or
6 tackifier, and water. As an alternative to commercial hydromulch, the
7 numerous other agricultural waste fibers, mulches and composts may be
8 utilized. Such may be preferred to favor the growth of certain species with
9 specialized requirements--for example, *Volvariella volvacea*, the Paddy Straw
10 mushroom, where rice straw is a preferred substrate. The fungal mycelium
11 which develops after application not only assists the growth of plants and
12 recovery of the ecosystem as above, but also serves to enhance the tenacity of
13 the fabric or geocloth, the many miles of mycelial hyphae forming widespread
14 connections between the cloth and the ground, thus preventing 'slippage' and
15 anchoring the fabric cloths, mulch, wood chips, straw, etc.

16 If desired, the hydroseeding mulch may optionally be partially
17 overgrown or completely overgrown with fungal mycelium prior to use. For
18 example, inoculation and growth for 48 to 72 hours will produce a
19 germinated, actively growing mycelium. Such mulches may be utilized with
20 fresh, actively growing mycelium or may be metabolically suspended via
21 refrigeration, drying or freeze-drying for storage and transport prior to

1 reactivation and use.

2 A wide variety of landscaping substrates, carriers, products and
3 materials are suitable for practice for the various embodiments of the present
4 invention. Where a bulk substrate mulch is desired, as for example in spray
5 hydroseeding of geocloths utilized to prevent erosion, suitable chopped,
6 chipped, shredded, ground, etc. fiber substrates include by way of example
7 (but not of limitation) woody and non-woody fibers such as wood chips,
8 sawdust, wood pulp, wood mulch, wood wastes, wood pellets and paper fiber
9 pellets, leaf paper, wood-based papers, non-wood papers, pressed cardboard
10 and corrugated cardboard, fiberized rag stock, cellophane, hemp and hemp-
11 like materials, bamboo, papyrus, jute, flax, sisal, coconut fibers and coir,
12 wheat straw, rice straw, rye straw, oat straw and other cereal straws, reeds,
13 rye grass and other grasses, grain hulls and other seed hulls such as
14 cottonseed hulls, cornstalks, corncobs or ground corncobs, soybean roughage,
15 coffee plants, waste and pulp, sugar cane bagasse, banana fronds, palm
16 leaves, the hulls of nuts such as almonds, walnuts, sunflower, pecans,
17 peanuts, etc., soy waste, cactus waste, tea leaves and a wide variety of other
18 agricultural waste products and combinations thereof. Suitable animal fibers
19 include wool, hair and hide (leather) and combinations thereof.

20 Alternatively, such pressurized spray hydroseeding may be utilized
21 without a cloth for landscaping, agriculture, covering garbage dumps (thus

1 preventing blowing garbage and dispersal by winds and ultimately enabling
2 improved biodegradation of dump materials) and numerous other
3 applications, with the water-fungus-hydromulch mixture being spread over
4 large areas. Such an approach may be preferred where it is desired to avoid
5 the expense of landscaping fabrics or geocloths, the time and effort of
6 installing and securing such fabric blankets, preparation of a relatively
7 smooth surface for installation, etc. The non-fungal component may be
8 varied in the ways known to those skilled in the art to favor the applied
9 fungal species, for example woodland mushrooms, grassland mushrooms,
10 dung inhabiting mushrooms, compost/litter/disturbed habitat mushrooms,
11 mycorrhizal mushrooms, entomopathogenic fungi and combinations thereof.

12 Using a subset of non-germinating seeds, and/or the outer shells and
13 hulls of germinating seeds within the propelled hydroseed mixture as food,
14 the mycelium can co-exist with germinating seeds in the applied
15 environment, benefiting both, and strengthening ecological fortitude.

16 Binding agents or “tackifiers” are typically preferably employed as a
17 component of the hydromulch. The tackifier/binding agent component of the
18 mulch enhances the strength and integrity of a mat-like tackified mulch
19 structure and may assist in adhering the mulch structure to the surface upon
20 which it is applied, assisting in the erosion control function and preventing
21 dispersal of the mulch from wind, rain, etc. Various binding agents and

1 tackifiers are known to those skilled in the art; see, for example, U.S. Patent
2 5,459,181 (1995) to West *et al.*

3 For many landscaping and agricultural applications, use of cart-
4 mounted hydroseeding units and the mobile hydroseeding variations will be
5 preferable. Such units are typically utilized to plant lawn grasses, and may
6 be utilized to plant native grasses, wildflowers, mixtures of grasses, shrubs,
7 bushes, trees, crops, etc. if desired. Spores, fresh mycelium, dried or freeze-
8 dried mycelium, powdered mushroom fruitbodies, the many forms of fungi
9 imperfecti and their conidia (asexually produced spores) and related fungal
10 forms and combinations thereof may be easily added to the hydroseeding
11 mixture. Hydroseeding units typically employ mechanical agitation (via
12 paddles or augers inside the tank) or jet mixing (via pump jets) of water and
13 materials; other methods will be readily apparent to those skilled in the art.

14 Hydroseeding as a fungal mycotechnology works well for numerous
15 reasons. The spores, mycelium or powdered mushroom fruitbodies and the
16 seeds are suspended in a nutrient rich slurry. The contact of the fungal
17 inoculum and seeds with the water triggers the germination cycle of both.
18 The mulch layer seals in the moisture and holds the soil in place (particularly
19 if a tackifier is utilized). The fungal inocula and seed are at an ideal depth
20 for good results. The conditions are right to produce lush growth in a very
21 short time. In addition, such an approach can greatly lower labor costs, with

1 one person simultaneously applying fungal inoculum, hydromulch, seed,
2 fertilizer and tackifier if desired, and water.

3 For use with trees and other slow germinating plants, a cover crop of,
4 for example, grass seeds or sterile hybrids can be applied in the mixture to
5 give a fast germinating ground cover, the grasses typically germinating first
6 followed by germination of the tree seeds. Alternatively, tree seedlings may
7 be directly utilized. As another example, a cover crop of millet or ryegrass or
8 sterile wheat can also be applied in the mixture to give a fast germinating
9 ground cover until the grass (or native grasses, etc.) being planted becomes
10 established. This method is only recommended for use during the growing
11 season of the particular grass species. Another preferred embodiment utilizes
12 a non-seeding annual grass, with the more expensive non-native grasses
13 being seeded at a later time after the nurturing biosystem has been
14 established.

15 Another preferred embodiment of the present invention is the use of
16 fungal inocula with agricultural equipment, including planting equipment,
17 harvesting equipment, field preparation equipment and processing
18 equipment with means for delivering fungal inocula. Appropriate methods of
19 modifying agricultural equipment with pumps, sprayers and/or mixers, etc. or
20 of mixing the fungal inocula with seeds (via the slurries above or other
21 means) will be readily apparent to those skilled in the art. Spores, mycelial

1 hyphae and or powdered mushrooms may be introduced into agricultural
2 equipment as liquids, powders, foams, sprays, creams, etc. and combinations
3 thereof or via other methods known to the art so as to provide the benefits of
4 simultaneous inoculation with saprophytic fungi, mycorrhizal fungi,
5 entomopathogenic fungi and/or other beneficial fungi. Alternatively, the
6 fungal inocula may be mixed with seeds and then distributed by the various
7 forms of agricultural planting equipment.

8 By way of example but not of limitation, such agricultural planting
9 equipment may include seeders, air seeders, planters, air planters, plate
10 planters, vacuum planters, drills, air drills, air seeding systems, row crop
11 cultivators, planting systems, inter-row or between row planting systems, rice
12 transplanters, etc.

13 Agricultural harvesting equipment may include, by way of example
14 only, combines, round balers, square balers, hay cubers, threshers and
15 threshing machines, forage harvesters, windrowers, rakes, tedders, mowers,
16 rotary mowers, sicklebar mowers, slashers and cutters, straw choppers, stalk
17 choppers, corn pickers, cotton strippers and gins, corn huskers, shellers, rice
18 harvesters, mechanical fruit and nut pickers, loaders, etc. The fungal inocula
19 may be utilized in various manners according to the desired purpose. For
20 example, it may be utilized to inoculate the remaining agricultural waste
21 and/or fields after harvest, thereby providing the numerous advantages

1 discussed herein via inoculation of the agricultural wastes and/or crop fields.
2 Alternatively, the fungal inocula may be utilized to directly inoculate the
3 agricultural products for uses as described herein, for example inoculation of
4 hay or straw with round or square balers, inoculation of hay with tedders,
5 inoculation of grasses with mowers, inoculation of corn husks and corn cobs
6 with huskers and shellers, inoculation of cotton wastes via cotton pickers and
7 strippers, inoculation of cotton seeds and hulls via cotton gins, inoculation via
8 loaders, etc.

9 In another preferred embodiment, such fungal inocula may be utilized
10 directly with agricultural equipment useful for preparation and/or
11 improvement of fields, orchards, etc. Such equipment includes by way of
12 example sprayers, irrigators, plows, cultivators, air carts, tillers and tillage
13 equipment, disks, openers, rippers, harrows, rotary hoes, blades, flail
14 shredders, flail cutters, rotary cutters, manure spreaders, flame weeders,
15 pruning machines, skids, scrapers, loaders, fertilizer spin spreaders,
16 pendulum spreaders, etc.

17 In another preferred embodiment, fungal spores and/or mycelium is
18 introduced into shredders and/or chippers to inoculate organic debris laid
19 onto landscapes.

20 The use of fungal inoculants as described above results in a
21 'mycofiltration' membrane lessening the impact of biological pathogens and

1 chemical pollutants in downstream environments. The fine network of
2 mycelial cells catches bacteria and other biological organisms as well as
3 releasing chemical agents (enzymes, peroxidases and acids) which decompose
4 toxins. In one field experiment, beds of *Stropharia rugosoannulata* were
5 established on dump truck loads of wood chips in ravines that drained from
6 pastures with a small herd of cattle onto a saltwater beach where oysters and
7 clams were being commercially cultivated. Prior to installing these beds,
8 fecal coliform bacteria seriously threatened the water quality. Once the
9 mycelium fully permeated the sawdust/wood chip beds, downstream fecal
10 bacteria were largely eliminated. The properly located mushroom beds
11 effectively filtered and cleaned the 'gray water' runoff of bacteria and
12 nitrogen-rich effluent. This observation was the stimulus for subsequent
13 study by Stamets, *Mycofiltration of gray water runoff utilizing Stropharia*
14 *rugosoannulata, a white rot fungus* (1993) (Unpublished Research Proposal
15 awarded a grant by the Mason County Water Conservation District, Shelton,
16 Washington). By using the fungal inoculation mycotechnologies disclosed
17 herein, such as 'mycocloths,' 'mycomats,' 'mycobags,' 'mycogabions' and
18 'mycoberms,' such results may be more efficiently and economically
19 accomplished. Such products and methods are in accord with the nature of
20 fungi--riparian habitat buffer zones work primarily because of mycelium.
21 Such colonized mycelial products will thus sequester nitrogen, carbon,

1 phosphorus and other compounds, a novel consequence of actively placing
2 such mycomaterials. Biodegradable mycoberms and similar structures may
3 be built repeatedly over time as an ongoing renewable process.

4 Such mycelial products are useful for combating virulent bacteria,
5 protists and protozoa, viruses, nematodes, rotifers, etc., for example *Escheria*
6 *coli*, *Bacillus subtilis*, malaria (e.g., *Plasmodium falciparum*), cholera (*Vibrio*
7 *cholerae*), anthrax (*Bacillus anthracis*), *Pfiesteria* (*Pfiesteria piscicida*), a
8 dinoflagellate causing toxic blooms which may assume numerous forms
9 during its lifetime, including a difficult-to-detect cyst stage, an amoeboid
10 stage, and a toxic vegetative stage, water-borne diseases and biological
11 warfare (BW) pathogenic species. Other harmful biological organisms that
12 can be digested and destroyed by fungal mycelia include nematodes, rotifers
13 and insect pests. Thus by infusing mycelium into cloths, rugs, blankets,
14 berms, hydroseeding mulches, soils, etc., targeted disease organisms such as
15 bacteria, fungi, viruses, protozoa, rotifers, amoebas and nematodes can be
16 effectively reduced, ameliorating the downstream impact as well as in
17 residence. Most or all fungi have antibacterial properties; fungi that are
18 preferred for use against bacteria include, for example, *Stropharia*
19 *rugosoannulata*, *Pleurotus* spp. and *Fomes fomentarius*. *F. fomentarius*, a
20 mushroom from the old growth forest, produced an army of crystalline
21 entities advancing in front of the growing mycelium, disintegrating when

1 they encountered *E. coli*, sending a chemical signal back to the mother
2 mycelium that, in turn, generated what appears to be a customized macro-
3 crystal which attracted the motile bacteria by the thousands, summarily
4 stunning them. The advancing mycelium then consumed the *E. coli*,
5 effectively eliminating them from the environment.

6 Such an approach may not only combat virulent organisms, but also
7 has the potential to provide fungal products which may be useful in
8 treatment or mitigation of the growth of such diseases. For example, a water
9 extract of *Polyporus umbellatus* mushrooms obtained from the present
10 inventor (available c/o Fungi Perfecti LLC, P.O. Box 7634, Olympia, WA
11 98507) were found to exhibit 100% inhibition of the growth of *Plasmodium*
12 *falciparum* during *in vitro* assays (Lovy *et al.*, Activity of Edible Mushrooms
13 Against the Growth of Human T4 Leukemic Cancer Cells, HeLa Cervical
14 Cancer Cells, and *Plasmodium falciparum*, *J. Herbs, Spices & Medicinal*
15 *Plants*, 6(4): 49-57 (1999)).

16 Toxic wastes, contaminants and pollutants that may be remediated by
17 the products and processes of the present invention include, by way of
18 example but not of limitation, organic compounds (taking advantage of the
19 unparalleled ability of fungi to degrade both naturally occurring and
20 synthetic organic molecules), inorganic compounds, and biological
21 contaminants including living organisms such as bacteria, viruses, protists,

1 nematodes, rotifers and combinations thereof.

2 More specifically, by way of example only, such organic compounds
3 include hydrocarbons such as polynuclear aromatic hydrocarbons (PAHs),
4 cyclic hydrocarbons and hydrocarbon chains such as alkanes and alkenes,
5 including the components of lubricants, fuels and solvents and additives such
6 as methyl t-butyl ether (MTBE), fertilizers, chemical pesticides including
7 organophosphate pesticides and organochlorines such as DDT
8 (dichlorodiphenyltrichloroethane), chlordane and toxaphene, the many
9 dioxins such as 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) and related
10 furans, organochlorines and organobromides such as pentachlorophenol
11 (PCP), polychlorinated biphenyls (PCBs) and polybrominated biphenyls
12 (PBBs), nitrogenous compounds such as such as ammonium nitrate, urea,
13 purines and putriscines, chemical warfare (CW) agents and nerve gases such
14 as the organophosphates Sarin (GB or O-isopropyl
15 methylphosphonofluoridate), Soman (GD or pinacolyl
16 methylphosphonofluoridate), Tabun (GA or O-ethyl N,N-
17 dimethylphosphoramidocyanidate), VX (O-ethyl S-[2-diisopropylaminoethyl]
18 methylphosphonothiolate) and VX family compounds, and their surrogates
19 such as isopropyl methylphosphonic acid (IMPA) and dimethyl
20 methylphosphonate (DMMP), and combinations thereof. One polypore
21 mushroom in the inventor's culture collection destroys the core constituent

1 base of the toxic nerve gas agents VX and Sarin. The fungi are also useful for
2 remediation of explosives (such as gunpowder and trinitrotoluene (TNT)),
3 explosive residues and explosives manufacturing byproducts (such as
4 dinitrotoluene (DNT)). By using cold-weather fungal strains, temperature-
5 sensitive munitions can be decomposed without the dangerous heat build-up
6 associated with typical compost mycoflora. Other contaminants that may be
7 remediated by the present invention include by way of example creosote,
8 alkaloids such as caffeine, endocrine-disrupting compounds such as estradiol,
9 steroids and other hormones, pro-hormones or hormone-like compounds,
10 detergents and soaps, textile dye pollutants including aromatic dyes, medical
11 wastes, urban runoff, industrial wastes and the many other toxic or
12 unpleasant byproducts of human activities. Such fungal products infused
13 with fungi capable of decomposing biological and chemical warfare toxins and
14 industrial toxins can be used to decontaminate toxic landscapes, battlefield
15 and otherwise, thus leading to reuse of valuable land.

16 One preferred type of fungal blanket, mat, bag or gabion is designed
17 specifically to treat oil spills and slicks. The mycomaterial is preferably made
18 of adsorbent biodegradable fiber materials and inoculated with spores and/or
19 hyphae of oil-eating fungi. Thus the oil is soaked up by the mat material and
20 digested by the mycelium of the fungus. A strain of *Pleurotus ostreatus* has
21 proven particularly effective in digesting and breaking down petroleum oils

1 (PAHs and alkanes); other preferred species include, by way of example but
2 not of limitation, *Trametes versicolor*, *Ganoderma lucidum* and other fungal
3 species as listed below. For soaking up and bioremediating spills on ocean
4 beaches, salt-water marsh fungi are typically preferable, for example
5 *Psilocybe azurescens*, *Psilocybe cyanescens* and *Flavodon flavus*.

6 Phosphorylated compounds such as the chemical warfare gases and
7 many organophosphate pesticides have proven particularly resistant to
8 breakdown and bioremediation, as few organisms are equipped with the
9 appropriate dephosphorylating enzymes. Fungi, on the other hand, have a
10 number of enzyme systems and paths for dealing with phosphorylated
11 compounds and are therefore particularly suited for remediation of
12 organophosphates. Preferred species include polypore fungi such as *Trametes*
13 *versicolor*, *Fomes fomentarius*, *Fomitopsis officinalis*, *Fomitopsis pinicola*,
14 *Phellinus igniarius*, *Phellinus linteus* and the other polypores listed below,
15 agarics such as *Psilocybe azurescens* and *Psilocybe cyanescens* containing
16 phosphorylated tryptamine compounds and their dephosphorylated analogs,
17 luminescent fungi utilizing adenosine triphosphate, luciferin and luciferase
18 for bioluminescence, and other phosphorus-rich mushroom fungi such as
19 *Agrocybe arvalis*, *Collybia* (*C. tuberosa* and *C. aluminosa*), *Coprinus*
20 *comatus*, *Lycoperdon perlatum* and *L. lilacinum*, *Pleurotus* species, esp. *P.*
21 *ostreatus* and *P. tuberregium* and *Psathyrella*, i.e. *P. hydrophila*.

1 Combinations may be preferred in certain applications as bringing a broad
2 range of phosphorus related enzymes to bear.

3 Since both *Psilocybe azurescens* and *Psilocybe cyanescens* can possess
4 up to 1-2% psilocybin, a phosphorus rich molecule, and/or psilocin, the
5 product of dephosphorylation of psilocybin, these species can be used to
6 dephosphorylate toxins wherein phosphorus contributes to the toxicity of the
7 pollutant (such as the phosphorylated chemical warfare gases above and
8 organophosphate pesticides). Grassland species such as *Psilocybe*
9 *semilanceata*, also rich in psilocybin, may also be preferably employed; such
10 grassland species have the advantageous characteristic of acting as
11 saprophytes, decomposing organic matter, or acting as ectomycorrhizal
12 species, directly benefiting plants via symbiosis, depending upon
13 circumstances. The non-psilocybin producing Blue *Stropharia* (blue-staining)
14 species can also be phosphorus containing and equipped with
15 dephosphorylating enzymes. These species include *Stropharia aeruginosa*, *S.*
16 *cyanea*, *S. albocyanea* and *S. caerulea*, and may be substituted where laws
17 restrict the use of the psilocybin-positive species, as may non-psilocybin
18 containing blue-staining *Panaeolus*, *Conocybe*, *Gymnopilus*, *Inocybe* and
19 *Pluteus*. Alternatively, specific enzyme blockers and/or other agents that
20 block the biosynthetic pathway of psilocybin and psilocin may be utilized. In
21 another approach, the *Psilocybe* species, which are known to take up

1 substituted tryptamines and convert them to non-naturally occurring analogs
2 of the natural tryptamine products, may be fed a substituted tryptamine that
3 would, on 4-hydroxylation or phosphorylation, produce an inactive compound.
4 Such substitution may be in the 4- position or in the 2-, 5-, 6-, N-, *alpha*-, etc.
5 positions or combinations thereof. Such substituted tryptamine analogs may
6 thus block or overwhelm the natural enzymes and phosphorus compounds.
7 Similarly, the phosphates such as organophosphate pesticides or nerve gases
8 may be used to overwhelm the naturally occurring enzymes to the exclusion
9 of naturally occurring psilocybin and psilocin. As another alternative, non-
10 fruiting strains of *Psilocybe* may be selected. As yet another alternative,
11 *Psilocybe* strains may be used solely in a mycelial state prior to the
12 production of psilocybin and psilocin--for example, it has been found with
13 *Psilocybe cyanescens* that no psilocybin or psilocin is formed in pre-primordial
14 mycelium, the mycelium knot stage of the mushroom being the earliest stage
15 at which psychoactive compounds could be detected. Gross, *J. Forensic Sci.*,
16 45(3): 527-37 (May 2000).

17 Luminescent mushrooms such as *Armillaria mellea*, *A. gallica*, *A.*
18 *bulbosa*, *Mycena citricolor*, *M. chlorophos*, *Omphalotus olearius* (*Clitocybe*
19 *illudens*) and *Panellus stypticus* present another example pathway of
20 phosphorus utilization by fungi that may be combined with the non-
21 luminescent species. Like the firefly and other organisms, fungi may exhibit

1 bioluminescence involving enzymatic excitation of a molecule to a high-
2 energy state and return to a ground state, accompanied by the emission of
3 visible light. Important molecular components are luciferin, a heat-stable
4 heterocyclic phenol and luciferase, a heat-labile enzyme. Luciferin and ATP
5 are thought to react on the catalytic site of luciferase to form luciferyl
6 adenylate, which is oxidized by molecular oxygen to yield oxyluciferin, which
7 emits light on returning to the ground state. A peroxide is presumed to be
8 formed as an intermediate.

9 The growth of algae in ponds and lakes can be directly attributed to
10 the phosphorus-rich runoff from agricultural fertilizers and other industrial
11 pollutants. Phosphorus is typically the 'limiting nutrient' of algae growth.
12 By removing phosphorus using mycocloths, mycomats and mycoberms
13 infused or spray hydroseeded with dephosphorylating fungi such as *Trametes*
14 *versicolor*, *Psilocybe azurescens*, and others, the over-growth algae can be
15 limited in lakes and ponds, providing cost and ecological saving benefits to
16 fishery ecologies and the watershed. A similar approach may be employed in
17 those soils and waters contaminated with organophosphate pesticide
18 residues. Floating mats of biodegradable materials may be infused with the
19 mycelia of anti-microbial fungi such as *Fomes fomentarius*, *Fomitopsis*
20 *officinalis*, *Ganoderma applanatum*, *Ganoderma oregonense*, *Trametes*
21 *versicolor*, *Lentinula edodes*, *Laetiporus sulphureus*, *Pleurotus eryngii*,

1 *Pleurotus ostreatus*, *Polyporus umbellatus*, *Psilocybe semilanceata*,
2 *Schizophyllum commune*, *Stropharia rugoso-annulata*, and *Calvatia* species
3 and placed into aquatic systems such as, but not limited to, ponds, lakes,
4 streams, rivers, and ditches for an effective treatment in reducing waterborne
5 disease microbes including but not limited to *Escherichia coli*, *Plasmodium*
6 *falciparum*, *Streptococcus* spp., *Staphylococcus* spp., *Listeria* spp., *Yersinia*
7 spp., *Shigella* spp.) and parasites (e.g., *Giardia* spp.)

8 Inorganic contaminants that may be remediated by fungi include by
9 way of example metals, phosphates, sulfates, nitrates, radionuclides and
10 combinations thereof. The fungal mycelia may or may not be able to
11 chemically alter an inorganic contaminant, for example metals or
12 radionuclides. However, the inorganic contaminant may be concentrated
13 from the surrounding ecological environment into fruiting bodies of the fungi.
14 With mixed organic/inorganic contaminants such as organometallic
15 compounds, the fungi may both degrade the compound and concentrate the
16 metal component.

17 The ability of higher fungi to concentrate heavy metals, metabolize
18 phosphorus compounds, etc., combined with the novel fiber products and
19 methods of the present invention allows use of fungally impregnated
20 materials, within or in absence of a matrix of biodegradable or non-
21 biodegradable materials, to sequester and concentrate heavy metals,

1 radioactive or otherwise, which then can be removed to eliminate toxins
2 topically and subsurface. Metallic effluents and ores may be treated with
3 specifically targeted fungi, for example the phosphate remediating
4 mushrooms for phosphate ores and runoff and/or metal concentrating
5 mushroom fungi. In addition, the fungi may favorably metabolize the organic
6 portion of organometallic compounds via mycofiltration and
7 mycoremediation.

8 Such residual organic debris from mycelia and the delivery systems
9 herein could be economically or profitably separated from the metals through
10 incineration, biodigestion with other organisms such as bacteria, protozoa,
11 yeasts, and/or via chemical treatments including acids, enzymes and
12 catalysts, including also the many other approaches known to the art. Such
13 an approach can also be favorably employed to control metal-laden runoff
14 from gold mines, silver mines, uranium mines, etc., providing control of mine
15 wastes while concentrating the valuable residual metals. Once sequestered
16 and concentrated, the metals may be removed by mechanical, chemical and/or
17 biological means. A number of mushroom fungi are known to concentrate
18 metals, including various edible mushrooms. One family of preferred genera
19 is *Collybia* and the similar *Marasmius* and their numerous "satellite genera"
20 in this 'taxonomically troubled' group. Such satellite genera (*Collybia* 'sensu
21 lato') include *Caulorhiza*, *Oudemansiella*, *Flammulina*, *Crinipellis*,

1 *Callistosporium*, *Micromphale* and *Marassmiellus*.

2 Examples of previous methodologies include those disclosed in U.S.
3 Patent No. 5,021,088 (1991) to Portier for separation and recovery of gold and
4 U.S. Patent No. 4,732,681 (1988) to Galun *et al.* for methods and systems for
5 use of a strain of *Cladosporium cladosporioides* to decrease heavy metal
6 concentrations such as lead, zinc, cadmium, nickel, copper and chromium in
7 industrial effluents. These and other similar methods may optionally be
8 combined with the higher fungi and the present invention for improved
9 separation and recovery from carbonaceous or pyritic or phosphate ores and
10 combinations thereof, including both gold and non-gold heavy metals such as
11 the radioactive and toxic metals. Thus the ore or industrial effluents
12 containing the various heavy metals may be treated with microorganisms,
13 such as fungi imperfecti and/or autotrophic bacteria such as *Thiobacillus*
14 *ferrooxidans* and *T. thlooxidans*, to leach soluble iron, copper and other metals
15 and sulfuric acid via oxidation of iron and sulfur prior to treatment with the
16 delivery systems of the present invention.

17 U.S. Patent No. 4,021,368 (1977) to Nemec *et al.* discloses use of lower
18 fungi microorganisms combined with polymers to “stiffen” the fungus and
19 eliminate the typical problems arising from fungi in general having a low
20 long term mechanical rigidity, causing difficulties in retention or absorption.
21 A stiff, coherent mycelial mat as provided by the delivery systems of the

1 present invention would be advantageous for collection of metal-enriched
2 mycelium and or mushrooms. Such may be provided via the present
3 invention in the form of a landscaping blanket, rug or mat or via bags or
4 gabions or via hydroseed fungal inoculation, optionally reinforced by a
5 polymer, metal or biodegradable fiber or combination thereof or other
6 support, with or without barrier materials ranging from tarps to complex
7 barriers . Alternatively, such supports and/or barriers may be utilized with
8 spray hydroseeding of hydromulch, wood chips, straw, etc., optionally with
9 tackifier, with 'sandwich' inoculation if desired, with or without fiber cloths or
10 gabions or such, so that the fungal species form a coherent, matlike
11 mycelium. Such an approach is also useful for biological concentration of
12 ores, ore slurries, etc., particularly of the heavy metals, as well as the various
13 other applications disclosed herein for mycoremediation, mycofiltration,
14 mushroom and plant cultivation, etc.

15 With or without such treatment with lower fungi and/or bacteria, mine
16 waste, effluent or ore substrate can be inoculated with saprophytic
17 mushrooms known for high yields, thereby allowing for the further
18 concentrating and sequestering of precious metals, toxic metals such as lead,
19 and/or the radioactive metals, both toxic and precious. For instance, Oyster
20 mushrooms, *Pleurotus ostreatus*, commonly convert 10% of the dry mass of
21 the substrate into dried mushrooms, allowing for a 'harvested' crop which can

1 be efficiently removed from the background environment. Subsequent to
2 Oyster mushrooms ceasing flushes, another species of mushrooms can be
3 introduced, such as *Stropharia rugoso-annulata*, which can further
4 concentrate the targeted compounds. Another round of concentration may be
5 carried out at that point by the numerous mushrooms which will grow upon
6 the rich soil that has been created via lignin degradation, including
7 mushrooms such as the 'Shaggy Mane,' *Coprinus comatus*, and the wide
8 variety of mushroom species ranging from gourmet lawn and field
9 mushrooms to little brown mushrooms to 'poisonous to humans' mushrooms.
10 By sequencing accumulator and hyperaccumulator mushroom species,
11 progressively greater extraction and/or concentration of valuable metals can
12 be achieved.

13 The fungal delivery systems of the present invention may also be
14 favorably combined with the techniques of phytoremediation (bioremediation
15 via plants) for maximum effectiveness of bioremediation of metals, persistent
16 organics, chlorinated organics, organophosphates, etc., including those 400+
17 plants that have to date been found to be "hyperaccumulators" of metals,
18 chlorinated solvents, etc. Suitable phytoremediation techniques for optional
19 combination with the delivery systems of the present invention include
20 phytoextraction (phytoaccumulation), rhizofiltration, phytostabilization,
21 phytodegradation (phytotransformation), rhizodegradation (enhanced

1 rhizosphere biodegradation), phytostimulation, or planted-assisted
2 bioremediation/degradation), and phytovolatilization. It is thought by the
3 present inventor and others that fungi assist and enable successful and
4 efficient hyperaccumulation via various direct and symbiotic mechanisms.

5 The present inventor has observed that one such preferred
6 hyperaccumulator species, the hybrid poplar, does particularly well in the
7 presence of saprophytic, wood decomposing mushrooms on wood chips and
8 fibrous media placed above the soil. By way of example only,
9 hyperaccumulator species for organics include poplars, cottonwood, mulberry,
10 juniper, sunflowers, fescues, ryegrasses and other grasses, clover, Indian
11 mustard, duckweed, parrotfeather, etc. and combinations of these and the
12 numerous other hyperaccumulators and accumulators found in the plant
13 world. Such hyperaccumulator species are, by way of example only, able to
14 extract and detoxify chlorinated solvent such as methylene chloride and
15 trichloroethylene (a major groundwater pollutant) and trinitrotoluene (TNT)
16 via the phytoremediation mechanisms as well as providing the known
17 admirable habitat improvement properties of healthy trees and plants via
18 shade, shelter, humidity maintenance, provision of lignin for conversion by
19 fungi into nutrients, etc.

20 In a preferred embodiment, poplars and other hyperaccumulator trees,
21 in symbiosis with fungi, display and maintain hydraulic control--mature

1 poplars have been estimated to transpire between 50 and 300 gallons of
2 water per day out of the ground. Hydraulic control is the use of plants to
3 rapidly uptake large volumes of water to contain or control the migration of
4 subsurface water. The water consumption by the poplars and other trees
5 decreases the tendency of surface contaminants to move towards ground
6 water and into drinking water. There are several applications that use
7 plants for this purpose, such as 'riparian corridors' or 'buffer strips' and
8 'vegetative caps.' Banks of poplars have also been used to stabilize
9 petroleum-contaminated groundwater flow, since the tree's prodigious
10 transpiration rate prevents movement of groundwater off site. The same
11 poplar technique has been shown to be an effective way to keep agricultural
12 runoff from entering streams, lowering pesticide and fertilizer contamination
13 of waterways, and thus may be favorably and advantageously combined with
14 the delivery systems and mycofiltration techniques of the present invention
15 which are separately able to perform large scale mycofiltration and
16 mycoremediation.

17 Hyperaccumulator plants are known in the scientific research and
18 patent literature that can concentrate metals thousands of times above
19 normal levels and can optionally be combined with the fungal delivery
20 systems for mine effluents and metallic ores described herein. For example,
21 planted on soil laden with nickel, *Streptanthus polygaloides* of the cabbage

1 family accumulates nickel up to one percent of its dry weight in its leaves and
2 flowers. Detoxifying the soil is as simple as harvesting the plants. The
3 'brake fern' (*Pteris vittata*) hyperaccumulates arsenic from contaminated soil,
4 attaining concentrations of arsenic as much as 200 times higher in the fern
5 than the concentrations in contaminated soils where it was growing. It will
6 accumulate arsenic even from soils having normal background arsenic levels.
7 As another example, after concentration and chelation via addition of a
8 chelating agent (or chelation and subsequent biological availability by the
9 present invention), lead can be accumulated by Indian mustard (*Brassica*
10 *juncea*). Indian mustard, in addition to lead, will hyperaccumulate
11 chromium, cadmium, nickel, selenium, zinc, copper, cesium, and strontium.
12 Sunflowers are known to absorb radioactive cesium and strontium, although
13 much of the metal remains bound in the root system, making it a poor
14 candidate for soil cleanup. After the 1986 Chernobyl nuclear disaster, Ilya
15 Raskin suspended sunflowers from Styrofoam rafts in ponds, where they
16 thrived, concentrating the metals up to 8,000 times the level in the water
17 itself, removing between 90 and 95 percent of the radioactivity from the pond.
18 The plants are removed, dried, and disposed of as radioactive waste. In
19 combination with the delivery systems of the present invention,
20 hyperaccumulators may optionally be employed with the fungal keystone
21 species, organic and inorganic nutrient gathering fungal species, and/or metal

1 concentrating fungal species and delivery systems of the present invention.

2 Whereas the literature of phytoremediation often teaches away from
3 use of fungi with plants or teaches the use of nutrient poor or nutrient limited
4 soils for some applications, often leading to poor hyperaccumulator growth,
5 such will typically not be the case when practiced with the present invention,
6 with or without added plant hyperaccumulators, as the fungi introduced by
7 the delivery systems herein tend to function as keystone species, leading to
8 lush habitats and vigorous growth of all plants, including hyperaccumulators,
9 with ecosystems better able to function as bioremediation agents.

10 Such fungally colonized mycelial products protect sensitive watersheds
11 such as salmon spawning grounds, providing mushroom and mycelial
12 biomass which then feed developing larvae of numerous insects which benefit
13 fisheries through enhancement of the food chain and from protection from
14 upland runoff. The present invention provides further advantages in
15 providing mycofiltration of pesticides, including both organophosphate and
16 halogenated pesticides, which are thought in minute quantities to interfere
17 with salmon's olfactory sense, thereby impeding the return to breeding
18 grounds and successful reproduction. Also provided are the sediment and silt
19 filtering advantages of mycofiltration. Sediment and silt runoff into salmon
20 and trout spawning grounds are known to create environment hostile to egg
21 survival. Similar negative habitat effects result from runoff into other bodies

1 of water. By utilizing mycofiltration, the silt and sediment becomes part of a
2 rich soil as opposed to a marine pollutant. The present invention as described
3 herein may be effectively employed to reduce, ameliorate, limit or prevent the
4 impact of pesticides and other agricultural and/or urban contaminants upon
5 riparian habitats and marine environments and the associated fisheries,
6 recreational use, drinking water, etc.

7 Fungi also present novel advantages in sequestration of carbon. The
8 international Kyoto Accords of 1998 helped establish a carbon-credit system,
9 an incentive-based system wherein those countries sequestering carbon,
10 effectively reducing the release of carbon dioxide, are rewarded. The concern
11 is to lessen the 'greenhouse effect', a major factor in global warming.

12 The no-till method of farming, wherein stubble is left for natural
13 decomposition, sequesters carbon in the soil. A study by Hu *et al.*, "Nitrogen
14 limitation of microbial decomposition in a grassland under elevated CO₂,"
15 *Nature*, 409: 188-191 (11 Jan. 2001), shows that elevation of carbon dioxide
16 levels in grasslands reduces microbial activity, specifically as seen through
17 the metabolism of nitrogen. Hence as CO₂ goes up, microbial activity goes
18 down. What these and other researchers have not yet recognized is that the
19 mycelium can intelligently regulate their grow-rates and out-gassing to
20 normalize the gaseous environment of the ecosystem in which they grow.
21 The cellular architecture of the fungal mycelial networks is made of carbon-

1 heavy molecules (chitin, carbohydrates and polysaccharides) and hence
2 habitats infused with mycelium using the present invention significantly
3 enhance their value in terms of augmented carbon credits.

4 In actively restoring devastated habitats using fungally impregnated
5 biodegradable materials, the current invention relies on the naturally gas-
6 governing properties of the selected fungal species. Encouraging the growth
7 of mycelium, and selecting the constellation of fungal species target-specific
8 to the toxic or threatened landscapes, enormous amounts of carbon can be
9 sequestered by the exoskeleton of the mycelial network, heavy in carbon-rich
10 molecules such as chitin and polysaccharides, and/or through the protein-rich
11 contents of the internal cell components. Furthermore, the active placement
12 of mycelial mosaics in a habitat additionally sequesters carbon directly
13 external to its cellular architecture through the production of extracellular
14 enzymes which convert cellulose precursor compounds into arabinoxylenes
15 and arabinogalactans. Mycelial mats of saprophytic and other fungi may
16 cover areas ranging from small plots to thousands of acres. The mushroom
17 mycelial mat is in fact a carbon bank.

18 The carbon credit system can also be economically applied when
19 incorporating the use of mycelium into organic debris fields and mycomats in
20 the reclamation of roads back into native ecosystems, optionally applying the
21 phytoremediation approaches above. Thousands of miles of roads must be

1 returned to natural conditions and the current energy crisis has caused 'hog
2 fuel' (= chipped junk wood used for furnaces) to skyrocket. The loss of carbon
3 from the ecosystem is an unfair economic practice as the hog fuel prices are
4 not being valued for their inherent carbon value. As governments
5 incorporate/recognize that the value of wood debris also should be considered
6 in terms of carbon credits, then the cost of using mycomats can be justified as
7 an economically valuable, cost-effective product and procedure for
8 incorporating carbon dioxide into fungi and plants in both microsphere and
9 biosphere.

10 Hence a major advantage of this invention is the active prevention of
11 atmospheric carbon dioxide through sequestering of carbon into the mycelial
12 network within the soil matrix. Thus, fungal growth can 'bank-roll' the
13 carbon credit system through such examples as the 'no-till' method and/or
14 through repairing threatened ecosystems by designing the insertion of
15 keystone fungi most beneficial to targeted environmental goals. By
16 sequestering carbon and increasing the value of the carbon credit, the
17 mycotechnologies of the present invention provide not only a cost effective
18 method, but also the numerous advantages arising from habitat
19 improvement.

20 Such landscaping substrates, cloths, carrier products, hydroseeding
21 equipment and agricultural equipment also provide means of introducing

1 mycorrhizal fungi. Such mycotechnologies also provide means for
2 introduction and “companion cultivation of saprophytic mushrooms” with
3 agricultural crops. The benefits of mycorrhizal fungi are well known; the
4 present inventor and others have also found that companion cultivation of
5 saprophytes enhances both quantity and quality of yields of grains and
6 vegetables and other crops. As mycelia bind soil particles (aggregation), soil
7 compaction is decreased and aeration is increased, allowing roots, oxygen,
8 carbon dioxide and water to move through the soil. This improvement in soil
9 quality may be noticed as a ‘bounce factor’ when walking over soils inoculated
10 with saprophytic fungi. For example, *Hypsizygus ulmarius* on sawdust,
11 covered with straw, has been found to be of great benefit to many crops and
12 plants, including corn, beans and Brussels sprouts; large ears of corn were
13 produced in a poor experimental soil, whereas previously the present inventor
14 had not been able to successfully cultivate corn in his garden due to growing
15 season and climate limitations. *Hypholoma sublateritium* was also of great
16 benefit to corn cultivation. *Stropharia rugosoannulata* is known to benefit
17 corn and was found to provide such a benefit, particularly in the second and
18 following years after inoculation. Thus companion cultivation of saprophytes
19 also offers preferred methods of improving crop yield while reducing the need
20 for fertilizers. See Pischl, C., *Die Auswirkungen von Pflanzen-*
21 *Pilzmischkulturen auf den Bodennaehrstoffgehalt und die Ernteertraege*

(1999), Master's Thesis, Leopold-Franzens-Universitat Innsbruck.
Mushrooms were observed fruiting underneath seedlings, the dewdrop formation and drip zone providing a preferred fruiting site. However, the plants and mushroom species must be carefully matched: while the Oyster-like mushroom *Hypsizygus ulmarius* had a beneficial effect on some neighboring crop plants, the Oyster mushroom *Pleurotus ostreatus* did not (Pischl, 1999). On the other hand, for nematode infested soils, *P. ostreatus* and other *Pleurotus* species may be preferred, the mycotechnologies herein acting as a nematode-control delivery system.

Inoculation of sawdust, straw or other fiber substrates placed on top of the soil has been found by the present inventor to be superior to and generally preferred to methods of inoculating and mixing with the soil for agricultural purposes; a more beneficial microclimate, microflora and biosphere results from placement of inoculated wood, straw, etc. on top of the soil. The no-till practice in particular improves the soil quality by fostering saprophyte populations that enhance the formation of water stable aggregates, thereby improving aeration, water infiltration, water retention and plant nutrient reserves. Such an approach also has the potential for producing gourmet and medicinal mushrooms.

The use of fungi (mycorrhizal and symbiotic saprophytic fungi) in a biodegradable matrix further aids the growth of resident and implanted flora.

1 Such examples include, but is not limited to the enhancement of native or
2 erosion-control grasses whose growth is enhanced from the fungal
3 components described herein. As the organic structural matrix, for example,
4 a straw/coconut cloth, is decomposed by the fungal component, grasses
5 benefit from the newly available nutrients liberated by the mycelium, from
6 the protective effect of the selected mycelium against invasive pathogenic
7 fungi and bacteria, and from the increase in water retention in otherwise
8 porous (sandy) soils. In both natural and man-made habitats, the
9 introduction of these fungi is an active component in enhancing
10 environmental health. For instance, the tenacity of *Ammophila maritima*, a
11 dune grass planted by the Army Corp of Engineers to prevent jetty erosion
12 around the Columbia River as it enters the Pacific Ocean, is significantly
13 enhanced through the domination of the mycelium of *Psilocybe azurescens*
14 and *P. cyanescens* in the top soils of that biosphere.

15 Of particular use where insect pest control is desired are the
16 entomopathogenic fungi *Metarhizium*, *Beauveria*, *Paecilomyces*, *Verticillium*,
17 *Hirsutella* and *Cordyceps*, either as the sole fungal species or in combination
18 with saprophytic and/or mycorrhizal species. In addition to known uses of
19 spores, the preconidial mycelium of entomopathogenic fungi has been found
20 to be attractant and/or pesticidal to such pest insects as termites, fire ants,
21 carpenter ants, etc. See U.S. patent application serial no. 09/678,141 (2000)

1 for MYCOPESTICIDES, U.S. patent application serial no. 09/922,361 (2001)
2 for MYCOATTRACTANTS AND MYCOPESTICIDES, and U.S. patent
3 application serial no. 09/969,456 (2001) for MYCOATTRACTANTS AND
4 MYCOPESTICIDES, all currently co-pending and herein incorporated in
5 their entirety by reference. Extracts of the pre-conidial mycelium of
6 entomopathogenic fungi, for example extracts of *Metarhizium*, *Beauveria*
7 and/or *Cordyceps*, are also useful for attracting and/or killing insects and may
8 be favorably combined with the fungal delivery systems disclosed herein. See
9 MYCOATTRACTANTS AND MYCOPESTICIDES above.

10 Insect pest control benefits are also provided by mycorrhizal fungi.
11 Plants infected by endophytic fungi are known to be chemically protected
12 against consumption by insect pests, for example aphids. Insect herbivore-
13 parasite interaction webs on endophyte-free grasses show enhanced insect
14 abundance at alternate trophic levels, higher rates of parasitism and
15 increased dominance by a few trophic links, whereas plants infected with
16 endophytes alter insect herbivore abundance, selectively favoring beneficial
17 insects and higher organisms. It is conceivable that the effect of plant
18 endosymbionts on food webs will cascade up through various trophic
19 pathways and can mediate competitive interactions between plant species
20 affecting vegetation diversity and succession. Ornacini *et. al.*, Symbiotic
21 fungal endophytes control insect host-parasite interaction webs, *Nature*, 409:

1 78-81 (4 January 2001). Thus in addition to their direct symbiotic effects
2 benefiting plants, it is expected that mycorrhizal fungi can reduce pest insect
3 herbivores, thus favoring beneficial insects and higher organisms and thereby
4 increasing biodiversity.

5 The parasitic fungi are particularly useful for the control and
6 extermination of invasive plant species, for example, the *Melaleuca* trees in
7 the Everglades. Such parasitic fungi include, for example, *Phellinus weirii*
8 and *Armillaria mellea*, two aggressive species. By use of non-sporulating
9 strains (as have been developed for *Pleurotus ostreatus*) incorporated into
10 myocloths or hydroseed spray, undesirable cross-infection outside of the
11 targeted area can be limited.

12 Control of plant pathogens such as *Rhizoctonia solani*, *Sclerotium*
13 *rolfsii*, *Verticillium dahliae* and other soilborne plant diseases may also be
14 provided by saprophytic and mycorrhizal fungi and by fungi imperfecti such
15 as *Trichoderma viride*, *T. harmatum* and *Gliocladium virens*.

16 Such mycotechnologies may be beneficial not only on Earth, but also
17 eventually in aiding the establishment of habitats in space colonies and in
18 the colonization of other planets. Such fabrics could be bio-engineered from
19 planetary surface dust ('soils') and impregnated with spores of fungi and
20 other organisms. Since there can be more than a billion spores per gram,
21 spores can be economically transported via drone or spaceship to the targeted

1 planetary body or space station. Their low weight/mass makes them
2 economically attractive bio-cargo for transportation through interplanetary
3 and interstellar space and the importance of fungi as a keystone species
4 makes them essential in any self-sustaining habitat.

5 Water and/or oils are preferably used to deliver spores and mycelial
6 hyphae, although spores and/or mycelium may be applied directly to the
7 landscaping materials, or traditional inoculation methods with grain and/or
8 sawdust spawn, etc. may be utilized (see Stamets, *Growing Gourmet and*
9 *Medicinal Mushrooms* (1993, 2000) and Stamets *et al.*, *The Mushroom*
10 *Cultivator* (1983), both herein incorporated by reference). Petroleum oils can
11 be readily digested by certain fungi (see U.S. patent application serial no.
12 09/259,077 (1999) for MYCOREMEDIATION (Thomas, Stamets *et al.*)),
13 currently co-pending, herein incorporated by reference) and biodegradable
14 oils are readily digested by most or all fungi perfecti and fungi imperfecti.
15 Therefore oil-spore or oil-hyphae mixtures or water-oil-spore or water-oil-
16 hyphae suspensions, with or without seeds, provide an alternative to the
17 water-spore or water-hyphae slurries which may be utilized in the practice of
18 the present invention. See also U.S. Patent Application Number 09/712,866
19 (2000) for SPORED OILS (Stamets), currently co-pending, herein
20 incorporated by reference. In general, where oils are utilized, biodegradable
21 oils are preferred as offering a more readily available nutritional source to a

1 wide variety of fungi. However, as some strains of white rot fungi have
2 proved to be voracious consumers of petroleum oils, species of oil-eating fungi
3 may be utilized with petroleum and mineral oil lubricants and synthetic and
4 semi-synthetic lubricants, as well as with biodegradable lubricants, vegetable
5 oil lubricants, modified vegetable oil lubricants, animal lubricants and
6 combinations and blends of these lubricants. Numerous vegetable oils are
7 suitable, including by way of example canola, rapeseed, castor, jojoba,
8 lesquerella, meadowfoam, safflower, sunflower, crambe, hemp, flax,
9 cottonseed, corn, olive, peanut, soybean and other such vegetable oil sources.
10 Such spored or hyphal oils may also be preferably employed in applications
11 such as ecological rehabilitation, mycoremediation and mushroom growing
12 where use of an oil as an additional nutritional source is desired.

13 The spores or fungal hyphae transfer agents may optionally contain
14 further amendments including germination enhancers, growth enhancers,
15 sugars, nutritional supplements, surface active and wetting agents, spore and
16 hyphae encapsulating materials, yeasts, bacteria, fungi imperfecti, etc.
17 Fungal hyphal mass can optionally be dried or freeze-dried and packaged,
18 with or without additional spores, in spoilage-proof containers for marketing
19 to end users as a seed and slurry additive. Fresh mycelial hyphae or mycelial
20 mass is best used immediately rather than stored for long periods.

21 Information on gathering useful and beneficial mushrooms for spores

1 or hyphae may be found in standard mycological field guides such as
2 *Mushrooms Demystified* (1979, 1986) by David Arora and *The Audubon*
3 *Society Field Guide to North American Mushrooms* (1981, 1995) by Gary
4 Lincoff.

5 As one gram of spores of, for example, *Ganoderma lucidum* may
6 contain more than a billion spores, it is therefore a simple matter to mix an
7 effective amount of spores into water or oil using mechanical or manual
8 mixing techniques known to the art and thereby provide a large number of
9 potential inoculation points.

10 Fungal spores may gathered via a variety of means, including but not
11 limited to large scale spore-printing on surfaces and collection from fresh
12 and/or dried mushrooms. A unique method developed by the present inventor
13 is to collect spores from the flexible poly-tubing or other ducting used for
14 distributing air within mushroom growing rooms and mushroom farms. This
15 method is efficient in gathering substantial spore mass.

16 Mycelial hyphae (including mushrooms, a form of mycelial hyphae)
17 may be cultured using standard mycological techniques for mushrooms.
18 Further information on techniques suitable for production of many of the
19 preferred gourmet, medicinal and eco-restorative mushrooms and their spores
20 and mycelial hyphae may be found in applicant's books, *Growing Gourmet*
21 *and Medicinal Mushrooms* and *The Mushroom Cultivator, supra*. One cost-

1 efficient method for expansion of mycelial mass for small-scale practice of the
2 present invention are commercial aerobic compost tea fermentors, which
3 allows growers to culture a very high concentration of aerobic
4 microorganisms in approximately 24 hours utilizing fine air particles infused
5 into the tea.

6 Virtually all fungi may be useful in habitat preservation and
7 restoration, reforestation and agriculture. Fungi useful in the present
8 invention include saprophytic fungi (including gilled, polypore and other
9 types of mushrooms), mycorrhizal fungi (which form a mutually dependent,
10 beneficial relationship with the roots of host plants ranging from trees to
11 grasses to agricultural crops, as may certain saprophytic fungi), and fungi
12 imperfecti (those asexually reproducing fungi related to the sexually
13 reproducing “fungi perfecti” or “mushroom fungi”). All fungi and their spores
14 and hyphae should be considered to be a useful part of the invention.

15 Suitable fungal genera include, by way of example but not of
16 limitation, the gilled mushrooms (Agaricales) *Agaricus*, *Agrocybe*, *Armillaria*,
17 *Clitocybe*, *Collybia*, *Conocybe*, *Coprinus*, *Flammulina*, *Giganopanus*,
18 *Gymnopilus*, *Hypholoma*, *Inocybe*, *Hypsizygus*, *Lentinula*, *Lentinus*, *Lenzites*,
19 *Lepiota*, *Lepista*, *Lyophyllum*, *Macrocybe*, *Marasmius*, *Mycena*, *Omphalotus*,
20 *Panaeolus*, *Panellus*, *Pholiota*, *Pleurotus*, *Pluteus*, *Psathyrella*, *Psilocybe*,
21 *Schizophyllum*, *Sparassis*, *Stropharia*, *Termitomyces*, *Tricholoma*,

1 *Volvariella*, etc.; the polypore mushrooms (Polyporaceae) *Albatrellus*,
2 *Antrodia*, *Bjerkandera*, *Bondarzewia*, *Bridgeoporus*, *Ceriporia*, *Coltricia*,
3 *Daedalea*, *Dentocorticium*, *Echinodontium*, *Fistulina*, *Flavodon*, *Fomes*,
4 *Fomitopsis*, *Ganoderma*, *Gloeophyllum*, *Grifola*, *Hericium*, *Heterobasidion*,
5 *Inonotus*, *Irpex*, *Laetiporus*, *Meripilus*, *Oligoporus*, *Oxyporus*, *Phaeolus*,
6 *Phellinus*, *Piptoporus*, *Polyporus*, *Schizopora*, *Trametes*, *Wolfiporia*, etc.;
7 Basidiomycetes such as *Auricularia*, *Calvatia*, *Ceriporiopsis*, *Coniophora*,
8 *Cyathus*, *Lycoperdon*, *Merulius*, *Phlebia*, *Serpula*, *Sparassis* and *Stereum*;
9 Ascomycetes such as *Cordyceps*, *Morchella*, *Tuber*, *Peziza*, etc.; 'jelly fungi'
10 such as *Tremella*; the mycorrhizal mushrooms (including both gilled and
11 polypore mushrooms) and endomycorrhizal and ectomycorrhizal non-
12 mushroom fungi such as *Acaulospora*, *Alpova*, *Amanita*, *Astraeus*, *Athelia*,
13 *Boletinellus*, *Boletus*, *Cantharellus*, *Cenococcum*, *Dentinum*, *Gigaspora*,
14 *Glomus*, *Gomphidius*, *Hebeloma*, *Lactarius*, *Paxillus*, *Piloderma*, *Pisolithus*,
15 *Rhizophagus*, *Rhizopogon*, *Rozites*, *Russula*, *Sclerocytis*, *Scleroderma*,
16 *Scutellospora*, *Suillus*, *Tuber*, etc.; fungi such as *Phanerochaete* (including
17 those such as *P. chrysosporium* with an imperfect state and *P. sordida*); the
18 fungi imperfecti and related molds and yeasts including *Actinomyces*,
19 *Alternaria*, *Aspergillus*, *Botrytis*, *Candida*, *Chaetomium*, *Chrysosporium*,
20 *Cladosporium*, *Cryptococcus*, *Dactylium*, *Doratomyces* (*Stysanus*),
21 *Epicoccum*, *Fusarium*, *Geotrichum*, *Gliocladium*, *Humicola*, *Monilia*, *Mucor*,

1 Mycelia Sterilia, *Mycogone*, *Neurospora*, *Papulospora*, *Penicillium*, *Rhizopus*,
2 *Scopulariopsis*, *Sepedonium*, *Streptomyces*, *Talaromyces*, *Torula*,
3 *Trichoderma*, *Trichothecium*, *Verticillium*, etc.; and entomopathogenic fungi
4 such as *Metarhizium*, *Beauveria*, *Paecilomyces*, *Verticillium*, *Hirsutella*,
5 *Aspergillus*, *Akanthomyces*, *Desmidiospora*, *Hymenostilbe*, *Mariannaea*,
6 *Nomuraea*, *Paraisaria*, *Tolypocladium*, *Spicaria*, *Botrytis*, *Rhizopus*, the
7 Entomophthoraceae and other Phycomycetes, and *Cordyceps*. It will be noted
8 that some entomopathogenic fungi imperfecti and molds can go through a
9 perfect stage, with the perfect form often getting a new name. It will also be
10 noted that such fungi imperfecti, molds and yeasts may produce spores,
11 conidia, perithecia, chlamydospores, etc. and other means of generating
12 progeny. All such fungi imperfecti, molds, yeasts, stages, forms and spores
13 should be considered as suitable for the practice of the present invention.

14 Suitable fungal species include by way of example only, but not of
15 limitation: *Agaricus augustus*, *A. blazei*, *A. brunnescens*, *A. campestris*, *A.*
16 *lilaceps*, *A. placomyces*, *A. subrufescens* and *A. sylvicola*, *Acaulospora delicata*;
17 *Agrocybe aegerita* and *A. arvalis*; *Albatrellus hirtus* and *A. syringae*; *Alpova*
18 *pachyploeus*; *Amanita muscaria*; *Antrodia carbonica*; *Armillaria bulbosa*, *A.*
19 *gallica*, *A. matsutake*, *A. mellea* and *A. ponderosa*; *Astraeus hygrometricus*;
20 *Athelia neuhoffii*; *Auricularia auricula* and *A. polytricha*; *Bjerkandera adusta*
21 and *B. adusta*; *Boletinellus merulioides*; *Boletus punctipes*; *Bondarzewia*

- 1 *berkeleyi*; *Bridgeoporus nobilissimus*; *Calvatia gigantea*; *Cenococcum*
- 2 *geophilum*; *Ceriporia purpurea*; *Ceriporiopsis subvermispora*; *Collybia*
- 3 *albuminosa* and *C. tuberosa*; *Coltricia perennis*; *Coniophora puteana*;
- 4 *Coprinus comatus* and 'Inky Caps'; *Cordyceps variabilis*, *C. facis*, *C.*
- 5 *subsessilis*, *C. myrmecophila*, *C. sphecocephala*, *C. entomorrhiza*, *C. gracilis*,
- 6 *C. militaris*, *C. washingtonensis*, *C. melolanthae*, *C. ravenelii*, *C. unilateralis*,
- 7 *C. clavulata* and *C. sinensis*; *Cyathus stercoreus*; *Daedalea quercina*;
- 8 *Dentocorticium sulphurellum*; *Echinodontium tinctorium*; *Fistulina hepatica*;
- 9 *Flammulina velutipes* and *F. populicola*; *Flavodon flavus*; *Fomes fomentarius*;
- 10 *Fomitopsis officinalis* and *F. pinicola*; *Ganoderma applanatum*, *G. australe*,
- 11 *G. curtisii*, *G. japonicum*, *G. lucidum*, *G. neo-japonicum*, *G. oregonense*, *G.*
- 12 *sinense* and *G. tsugae*; *Gigaspora gigantia*, *G. gilmorei*, *G. heterogama*, *G.*
- 13 *margarita*; *Gliocladium virens*; *Gloeophyllum saepearium*; *Glomus*
- 14 *aggregatum*, *G. caledonius*, *G. clarus*, *G. fasciculatum*, *G. fasciculatus*, *G.*
- 15 *lamellosum*, *G. macrocarpum* and *G. mosseae*; *Grifola frondosa*; *Hebeloma*
- 16 *anthracophilum* and *H. crustuliniforme*; *Hericiium abietes*, *H. coralloides*, *H.*
- 17 *erinaceus* and *H. capnoides*; *Heterobasidion annosum*; *Hypholoma capnoides*
- 18 and *H. sublateritium*; *Hypsizygus ulmarius* and *H. tessulatus* (= *H.*
- 19 *marmoreus*); *Inonotus hispidus* and *I. obliquus*; *Irpex lacteus*; *Lactarius*
- 20 *deliciosus*; *Laetiporus sulphureus* (= *Polyporus sulphureus*); *Lentinula edodes*;
- 21 *Lentinus lepideus*, *L. giganteus*, *L. ponderosa*, *L. squarrosulus* and *L.*

- 1 *tigrinus*; *Lentinula species*; *Lenzites betulina*; *Lepiota rachodes* and *L.*
- 2 *procera*; *Lepista nuda* (= *Clitocybe nuda*); *Lycoperdon lilacinum* and *L.*
- 3 *perlatum*; *Lyophyllum decastes*; *Macrocybe crassa*; *Marasmius oreades*;
- 4 *Meripilus giganteus*; *Merulius tremellosus* and *M. incarnatus*; *Morchella*
- 5 *angusticeps*, *M. crassipes* and *M. esculenta*; *Mycena citricolor* and *M.*
- 6 *chlorophos*; *Omphalotus olearius*; *Panellus stypticus*; *Paxillus involutus*;
- 7 *Penicillium oxalicium*; *Phaeolus schweinitzii*; *Phellinus igniarius* *P. linteus*
- 8 and *P. weirii*; *Pholiota nameko*; *Piloderma bicolor*; *Piptoporus betulinus*;
- 9 *Pisolithus tinctorius*; *Pleurotus citrinopileatus* (= *P. cornucopiae* var.
- 10 *citrinopileatus*), *P. cystidiosus*, (= *P. abalonus*, *P. smithii* (?)), *P. djamor* (= *P.*
- 11 *flabellatus*, *P. salmoneo-stramineus*), *P. dryinus*, *P. eryngii*, *P. euosmus*, *P.*
- 12 *ostreatus*, *P. pulmonarius* (= *P. sajor-caju*) and *P. tuberregium*; *Pluteus*
- 13 *cervinus*; *Polyporus indigenus*, *P. saporema*, *P. squamosus*, *P. tuberaster* and
- 14 *P. umbellatus* (= *Grifola umbellata*); *Psathyrella hydrophila*, *Psilocybe*
- 15 *aztecorum*, *P. azurescens*, *P. baeocystis*, *P. bohémica*, *P. caerulescens*, *P.*
- 16 *cubensis*, *P. cyanescens*, *P. hoogshagenii*, *P. mexicana*, *P. pelliculosa*, *P.*
- 17 *semilanceata*, *P. tampanensis* and *P. weilii*; *Rhizopogon nigrescens*, *R.*
- 18 *roseolus* and *R. tenuis* (= *Glomus tenuis*); *Schizophyllum commune*;
- 19 *Schizopora paradoxa*; *Sclerocytis sisuosa*; *Serpula lacrymans* and *S.*
- 20 *himantioides*; *Scleroderma albidum*, *S. aurantium* and *S. polyrhizum*;
- 21 *Scutellospora calospora*; *Sparassis crispa* and *S. herbstii*; *Stereum*

1 *complicatum* and *S. ostrea*; *Stropharia aeruginosa*, *S. cyanea*, *S. albocyanea*,
2 *S. caerulea* and *S. rugosoannulata*; *Suillus cothurnatus*; *Talaromyces flavus*;
3 *Termitomyces robustus*; *Trametes hirsuta*, *T. suaveolens* and *T. versicolor*;
4 *Trichoderma viride*, *T. harmatum*; *Tricholoma giganteum* and *T. magnivelare*
5 (Matsutake); *Tremella aurantia*, *T. fuciformis* and *T. mesenterica*; *Volvariella*
6 *volvacea*; and numerous other beneficial fungi.

7 For ecological restoration, all the fungi (including not only
8 economically valuable species but also “little brown mushrooms” and
9 “toadstools”) may play a valuable role, including stump and log dwelling
10 fungi, wood chip dwelling fungi, ground dwelling fungi, mycorrhizal fungi and
11 the fungi imperfecti. For example, spores or hyphae of the genus *Morchella*
12 such as *Morchella angusticeps*, *M. crassipes* and *M. esculenta*, gourmet
13 ground dwelling mushrooms that are known to favor fire-burned areas, may
14 optionally be utilized in the present inventions in fire recovery efforts,
15 thereby introducing a potential source of very rapidly growing mycelium into
16 the soil at the same time seeds are introduced or landscaping cloths are laid.
17 Preferred species for ecological restoration (and most other purposes) include
18 *Auricularia polytricha*; *Agaricus blazei* and *A. brunnescens*; *Agrocybe*
19 *aegerita*; *Bridgeoporus nobilissimus*; *Coprinus comatus*; *Flammulina velutipes*
20 and *F. populicola*; *Fomes fomentarius*; *Fomitopsis officinalis* and *F. pinicola*;
21 *Ganoderma lucidum*, *G. oregonense* and *G. tsugae*; *Grifola frondosa*; *Hericium*

1 *abietes* and *H. erinaceus*, *Hypholoma capnoides* and *H. sublateritium*;
2 *Hypsizygus ulmarius* and *H. tessulatus*; *Laetiporus sulphureus*; *Lentinula*
3 *edodes*; *Lepista nuda*; *Morchella angusticeps*; *Pholiota nameko*; *Pleurotus*
4 *citrinopileatus*, *P. cystidiosus*, *P. eryngii*, *P. euosmus*, *P. ostreatus*, *P.*
5 *pulmonarius* and *P. tuberregium*; *Polyporus umbellatus* and *P. tuberaster*;
6 *Psilocybe azurescens*, *P. cubensis*, *P. cyanescens*, *P. mexicana*, *P. semilanceata*
7 and *P. tampanensis* (where these species are legal for such purposes);
8 *Sparassis crispa*; *Stropharia rugosoannulata*; *Trametes versicolor*; *Tremella*
9 *fuciformis*; and *Volvariella volvacea*.

10 Of particular use where insect pest control is desired are the
11 entomopathogenic fungal species *Metarhizium anisopliae*, *Metarhizium*
12 *flaviride* , *Beauveria bassiana*, *Beauveria brongniartii*, *Beauveria amorpha*,
13 *Pacilomyces fumosoroseus*, *Verticillium lecanii*, *Hirsutella citriformis*,
14 *Hirsutella thompsoni*, *Cordyceps variabilis* , *Cordyceps facis*, *Cordyceps*
15 *subsessilis*, *Cordyceps myrmecophila*, *Cordyceps sphecocephala*, *Cordyceps*
16 *entomorrhiza*, *Cordyceps gracilis*, *Cordyceps militaris*, *Cordyceps*
17 *washingtonensis*, *Cordyceps melolanthae*, *Cordyceps ravenelii* , *Cordyceps*
18 *unilateralis* and *Cordyceps clavulata*.

19 Preferred species for mycoremediation include the saprophytic
20 mushrooms *Fomes fomentarius* (*E. Coli* and other bacteria, protists,
21 pathogens etc.); *Fomitopsis officinalis* and *F. pinicola*; *Ganoderma lucidum*,

1 *G. oregonense* and *G. tsugae*; *Laetiporus sulphureus*; *Pleurotus ostreatus* and
2 the other *Pleurotus* species (oils, polyaromatic, alkane and alkene
3 hydrocarbons including chlorinated compounds, brominated compounds,
4 hormones, etc.); *Polyporus umbellatus* (malaria and other bacteria); *Psilocybe*
5 *azureus* and *P. cyanescens* (Sarin and VX and other phosphorylated nerve
6 gases, organophosphate pesticides, etc.); *Stropharia rugosoannulata*
7 (bacteria, urban and agricultural runoff, mycofiltration, as a “follow-up”
8 species to *Pleurotus* and other white-rot fungi, etc.); and *Trametes versicolor*
9 and other *Trametes* and species (Sarin, VX and other phosphorylated nerve
10 gases, organophosphate pesticides, etc.), *Collybia* and the similar *Marasmius*
11 and numerous “satellite genera” (metals, heavy metals, ores, etc.) as well as
12 the other gilled and polypore genera and species listed above. Where the
13 mycotechnologies of the present invention are utilized for remediation of toxic
14 materials, the fungal species are preferably adapted to the substrate, that is
15 cultured, fed (challenged with) the target contaminant(s) or substrates,
16 selected for vigorous growth and thereby preconditioned to most effectively
17 degrade the target substrates and/or contaminant(s). See *Growing Gourmet*
18 *and Medicinal Mushrooms* and MYCOREMEDIATION, *supra*.

19 The species above include some of the many examples of the useful and
20 beneficial fungi that may be utilized with the present invention; the scope of
21 the invention as pertaining to fungi should not be considered thereby limited,

1 as it will be recognized that all fungi may be favorably employed in the
2 present invention.

3 By selecting the type of fungal spores or hyphae to be infused into the
4 target, the course of colonization by fungi can be directed, allowing selection
5 of economically or ecologically significant species of fungi, including
6 mushrooms useful for ecological preservation, reforestation and habitat
7 restoration, mushrooms useful for bioremediation of toxic wastes and
8 pollutants, mushrooms with mycelia useful as an agricultural amendment,
9 gourmet mushrooms, medicinal mushrooms containing valuable
10 physiologically active compounds and pro-compounds, and mushrooms
11 containing valuable enzymes, enzyme precursors and useful chemical
12 compounds. Succession also occurs--as one type of mushroom exhausts its
13 nutrient supply, another takes its place. To some degree, control of the
14 successions of insect populations can also be achieved by selecting mosaics of
15 fungal species which can predetermine species sequences. Fungal species
16 may be selected for a specific environment, for example lawns, gardens, crop
17 fields, forests (ranging from plains to mountainous to tropical ecosystems
18 environments), aquatic environments including riparian, marsh, wetlands,
19 estuaries, ponds, lakes, ditches, saline environments, etc.

20 A single species may be employed for a single application--for example,
21 a single saprophytic species on a fiber substrate in conjunction with a single

1 plant species such as *Hypsizygus ulmarius* on sawdust with corn. For typical
2 ecological restoration, mycoremediation of toxic wastes, habitat restoration
3 and preservation, etc., a plurality of species is preferred. The variety of
4 species produce different species specific enzymatic systems that break down
5 different chemicals and make these chemicals biologically available as
6 nutrients for the microsphere and the biosphere. An example can be seen in
7 the breakdown of a recalcitrant substrate--a hardwood such as ironwood, a
8 substrate containing high concentrations of the complex polyaromatic
9 cellulose carbohydrate compounds and the complex heterogeneous
10 polyaromatic polymer lignin. A succession of mushrooms may be grown on
11 the same wood, each species breaking down different compounds via different
12 enzymatic systems, thereby making the carbon, nitrogen, phosphorus,
13 hydrogen, etc. available as nutrients. To illustrate, a succession of gourmet
14 mushroom species may be grown on the same wood. For example, *Lentinula*
15 *edodes* (Shiitake) may be first grown on the wood, then *Pleurotus ostreatus*
16 (Oyster), then *Stropharia rugosoannulata* (King Stropharia, Garden Giant or
17 'Godzilla Mushrooms'), at which point the wood will have been transformed
18 into a rich soil, suitable for gourmet mushrooms such as *Coprinus comatus*
19 (Shaggy Mane). The same principle can be observed in nature where three or
20 four different mushroom species may be observed fruiting from the same
21 stump, each digesting a different woody compound and making the

1 compounds available to the biosphere in the form of mycelium and
2 mushrooms, or where different species of mushrooms may be observed
3 fruiting from the floor of the forest adjacent to each other. The saprophytic
4 mushrooms illustrated above also make such nutrients available to
5 mycorrhizal fungi, thus further enhancing the symbiotic relationship with
6 plants and resulting in greatly increased growth. Thus a plurality of fungal
7 strains and species is often preferred, including, for example, the various
8 saprophytic mushroom fungi and combinations of fungi including
9 saprophytic-entomopathogenic, saprophytic-mycorrhizal, saprophytic-
10 mycorrhizal-entomopathogenic, saprophytic-mycorrhizal-fungi imperfecti,
11 etc., optionally packaged separately or in combination with seeds, the various
12 fiber substrates, soils, etc.

13 It will be appreciated that many or all seeds or seedlings may be
14 preferably employed with the present invention. While the totality of plants
15 is too large to list, a few examples of native grass, sedge, rush and grass-like
16 seeds and cultivated seeds include *Agrostis exarata* (Spike Bentgrass),
17 *Ammophila arenaria* (European sand dune or beach grass), *Ammophila*
18 *breviligulata* (American beach grass), *Ammophila champlainensis* Seymour,
19 *Ammophila maritima*, *Beckmannia zyzigachne* (American Sloughgrass),
20 *Bromus carinatus* (California Brome), *Bromus vulgaris* (Columbia Brome),
21 *Carex densa* (Dense-Headed Sedge), *Carex feta* (Green-Sheathed Sedge),

1 *Carex leporina* (Harefoot Sedge), *Carex lenticularis* (= *C. kelloggii*) (Shore
2 Sedge), *Carex lyngbyel* (Lyngby Sedge), *Carex macrocephala* (Big Headed
3 Sedge), *Carex obnupta* (Slough Sedge), *Carex pansa* (Foredune Sedge), *Carex*
4 *unilateralis* (One-Sided Sedge), *Deschampsia caespitosa* (Tufted Hair Grass),
5 *Eleocharis palustis* (Creeping Spike rush), *Elymus glaucus* (Blue Wild Rye),
6 *Festuca idahoensis*- var. *roemerii* (Roemer's Fescue), *Festuca rubra* var.
7 *littoralis* (Shore Fescue), *Festuca subulata* (Bearded Fescue), *Glyceria elata*
8 (Tall Mannagrass), *Glyceria occidentalis* (Western Mannagrass), *Hordeum*
9 *brachyantherum* (Meadow Barley), *Juncus effusus* (Soft Rush), *Juncus patens*
10 (Spreading Rush), *Juncus tenuis* (Slender Rush), *Lozula campestris*
11 (Woodrush), *Phalaris arundinacea* (Reed Canary Grass), *Phalaris aquatica*,
12 *Phalaris tuberosa* (Staggers Grass), *Phalaris canariensis*, *Poa Macrantha*
13 (Dune Bluegrass), *ReGreen* (Sterile Hybrid Wheat), *Scirpus acutus* (Hardstem
14 Bullrush), *Scirpus americanus*, *Scirpus cyperinus*, *Scirpus maritimus*
15 (Seacoast Bullrush), *Scirpus microcarpus*, *Scirpus validus*, *Sparaganuim*
16 *eurycarpum* (Giant Burreed), *Triglochin maritimum* (Seaside Arrowgrass),
17 *Typha latifolia* (Cattail), *Alopecurus geniculatus*, *Carex pachystachya*, *Carex*
18 *stipata* (grass like), *Danthonia californica*, *Eleocharis ovata* (grass like),
19 *Glycaria grandis*, *Juncus acuminatus*, *Juncus bolanderi* and *Juncus*
20 *ensifolius* (Daggar leaf rush).

21 Example applications include: 1) Habitat recovery/reclamation:

1 'regreening' of roads, especially logging roads, important in lands returned to
2 wilderness or wildlife preserves and for prevention of sediment and silt runoff
3 into waterways from existing gravel roads, depleted environments, scarred or
4 biologically hostile environments, all typically lacking topsoils. For example,
5 a preferred method of restoration on top of gravel logging roads would be to
6 lay down a 2.5-10 cm. (1-4 inch) layer of mixed wood chips (i.e. hog fuel type
7 wood chips), broadcast saprophytic and mycorrhizal species either by free
8 hand, hydroseeding or via mycocloths or mycobags (or any combination
9 thereof or via other mycotechnologies discussed herein), grass seeds are
10 applied, and then chopped straw, twigs, etc. loosely overlaid over the top
11 surface to provide shade and moist air pockets. If a non-seeding, non-native
12 grass, is used the first year, the carbon cycle is begun, and as they mature,
13 decline and die, the newly available debris further fuels the carbon cycle. By
14 using a light infusion of native seeds and/or seeds or seedlings of shrubs and
15 trees, or by depending upon natural re-seeding from adjacent lands, this
16 method will stimulate the process of habitat restoration leading to a more
17 native environment. The process of soil generation is sped up by months,
18 releasing nutrients to benefit plants and other organisms. This process
19 creates topsoils and encourages biological recovery and complexity. The
20 mycelium retains sediments and silts washed from the gravel road,
21 incorporating them into topsoil while preventing release into waterways.

1 This is also useful as a method of accumulating carbon credits.; 2)
2 Mycofiltration: protection of sensitive watersheds and ecosystems from
3 upland or neighboring sources/vectors of contamination by capturing in the
4 mycelial network. This is critical for urban developments, protection of
5 salmon or trout streams, estuary environments, etc.; 3) Mycobags,
6 mycogabions, mycoclths and mycobags overlaying toxic waste fields:
7 penetration of mycelium to several inches is achieved, a year later,
8 decontaminated soil can be scooped up (now a value added product), and then
9 another layer of mycobags, mycogabions, etc. can be placed on top. This can
10 be done sequentially for the deep removal of toxins.; 4) Saprophytic,
11 mycorrhizal-saprophytic-entomopathogenic, saprophytic-entomopathogenic
12 and other fungally inoculated substrates for environmental and agricultural
13 enhancement and control of pest microorganisms and insects; 5) Soil
14 regeneration and reforestation via burlap bags inoculated with fungi and
15 layered over the ground with hybrid poplars planted 6-12 feet apart; 6) Deep
16 trenching wherein a narrow, deep ravine is filled with sawdust, woodchips,
17 straw and/or agricultural wastes and inoculated with mycelium; 7) Chicken
18 (and other animal) farms where waste exceeds the capacity to recycle,
19 resulting in phosphorus and nitrogen devastating the watershed.
20 Mycofiltration is achieved via creation of 'mycological parks' utilizing species
21 suited to the local environmental conditions and wastes/nutrient materials

1 for fungal growth). For example, in the southeastern United States,
2 *Pleurotus ostreatus* and *P. eryngii*, *Coprinus comatus* and
3 *Agaricus brunnescens*, *A. blazei* and *A. bitorquis* could be used for sheet
4 inoculation, covered with 5-15 cm. (2-6 inches) of chicken/sawdust waste.
5 Poplars, cottonwoods and other trees could be planted for hydraulic control
6 and protection of groundwater; 8) A cardboard insect monitoring station
7 utilizing mycoattractants such as extracts of pre-conidial mycelia and/or pre-
8 conidial mycelia of mycopesticidal, entomopathogenic fungi such as
9 *Metarhizium anisopliae*, *Beauveria bassiana*, *Paecilomyces* and *Cordyceps*
10 species. Since the targeted insects respond to and are drawn towards the loci
11 of the extracts, the extracts can be presented in a wide variety of ways and
12 still demonstrate attractancy. The insect myco-attractant may be saturated
13 into a wicking agent or membrane to slowly out-gas the attractant fragrance.
14 The surface area of the membrane or wick, its absorptive properties, its rate
15 of release of volatile attractants and the duration of wicking are all
16 influenced and easily altered according to the target insect and
17 environmental considerations. The monitoring station would then register
18 'hits' by registering by any means the numbers of visitations from the insects.
19 This sampling can be indispensable for recommending subsequent
20 treatments; 9) Empowering other insect treatment and control systems. The
21 soaking of mycoattractant extract onto cellulose, paper, cardboard, wood or

1 other biodegradable materials for a period of time and at a concentration to
2 be effective allows for construction of a biodegradable monitoring or kill
3 station. The insects, such as termites, fire ants and carpenters ants, enter
4 into a chamber where the mycoattractant is localized and then are trapped
5 and/or killed via ingestion of the material containing mycopesticidal extract.
6 Alternatively, the target insects are attracted to the monitoring station, trap
7 or to a close proximity where they are captured and/or killed via any insect
8 treatment or control means, including but not limited to the use of adhesives,
9 electricity, moving air, sprays, chemicals (toxins, growth regulators, for
10 instance), desiccants, cold temperatures, hot air, mechanical devices and
11 combinations thereof. Such monitors or traps can be useful to analyzing,
12 treating and solving the problems associated with invasive insects, and is
13 highly applicable to rural, agricultural, forested, urban and suburban
14 settings. 10) Controlling social insects such as fire ants, carpenter ants and
15 termites with the construction of monitoring and/or killing stations utilizing
16 extracts of the pre-conidial mycelia of mycopesticidal, entomopathogenic
17 fungi combined with pre-conidial mycelium of such fungi on a biodegradable
18 cellulosic material like wood, paper or cardboard. This combination of extract
19 and live mycelium has two advantages. The target insects are attracted to
20 the locus from which the fragrance of the extract emanates. As the mycelia
21 grows, it also outgases an attractant fragrance. The insect consumes the

1 extract-impregnated cellulose and also makes contact with fragments of
2 mycelia. As the insect travels, mycelia is spread. As the insect weakens with
3 illness, the mycelia becomes stronger. The insect is killed by both exposure to
4 the attractant but toxic extract and from infectious colonization by the
5 fungus. The time delay of exposure to death is an added advantage as it
6 allows the infected individuals to fully disperse through the affected region as
7 well as the nest without being sequestered and expunged from the colony; 11)
8 The use of mycoattractants derived from the extract of the mycelia of pre-
9 conidial, entomopathogenic, mycopesticidal fungi to place 'bait stations'
10 having these extracts in strategic locations to draw in insect plagues to a
11 single locus. Locust plagues could be diverted and drawn towards 55 gallon
12 drums hosting the mycoattractants wherein the insects could be trapped.
13 Mycelially based extracts of pre-conidial mycelium of entomopathogenic fungi
14 could be utilized to prevent plagues, herd insects to control points, avoiding
15 massive crop damage and economic devastation, and negating the need for
16 costly and toxic chemicals; 12) The use of mycoattractants derived from the
17 extract of the mycelia of pre-conidial, entomopathogenic, mycopesticidal fungi
18 to draw in beneficial insects whose predatory preferences include the plague
19 insect. For instance, a gardener could increase the number of lady bugs if
20 aphid infestations get out of control; and 13) The use of attractant emitters
21 using extracts of pre-conidial mycelium from mycopesticidal,

1 entomopathogenic fungi to attract pollinating insects to disadvantaged plants
2 by placing them in close proximity of the targeted plants. This invention will
3 be become increasingly important with the loss of sufficient populations of
4 insects which would otherwise naturally accomplish the task of pollination.

5 EXAMPLE 1

6 A coconut fiber door mat was pressure steam-sterilized in a
7 polypropylene bag at 1 kg/cm² (15 psi) for two hours, inoculated with rye
8 grain spawn, and the fungus allowed to overgrow the mat. Grass seeds were
9 added and the mat moved to an outdoor location. The mat was observed to
10 fruit *Pleurotus ostreatus* (Oyster) mushrooms and the seed was observed to
11 sprout and prosper. Birds were observed hunting for grass seed in the
12 mycomat; they appeared to prefer feeding from the fungal mat as compared
13 to feeding from a nearby (15 feet) bird feeder. The birds were observed to add
14 bird guano to the mat, thereby increasing the nutritional base and
15 introducing various organisms to the biological community.

16 EXAMPLE 2

17 Grain spawn of *Pleurotus ostreatus* was layered between straw-
18 coconut fiber mats steam-sterilized as above. Oyster mushrooms pushed
19 through the un-colonized upper layer of the straw-coconut fiber mat,
20 resulting in 'island fruitings' scattered over the mats with a heavy dusting of
21 spores dispersed around the mushrooms. These parents provided the means

1 for subsequent and more thorough colonization. This sandwich inoculation
2 provides an extremely efficient use of spawn, with sheet inoculation of thin
3 layer(s) of spawn producing a prodigious amount of spores and numerous
4 satellite colonies of inoculated substrate.

5 EXAMPLE 3

6 By introducing spores of *Stropharia rugosoannulata*, an edible
7 mushroom, into hydroseeding mulch materials, the receiving fabric material,
8 straw and wood chips soon colonized with mycelium. Plant growth was
9 enhanced, as well as water retention, and eventually edible mushrooms were
10 produced. Bees were attracted to the mycelium and fly larvae hatched from
11 the mushrooms along the stream bank, the larvae and resultant insects
12 providing a benefit to fish. In two years the wood chips had become rich soil.

13 The present invention utilizes the design and active insertion of
14 individual saprophytic, mycorrhizal, entomopathogenic, and parasitic fungal
15 species and mosaics of species to catalyze habitat recoveries from
16 catastrophes. Furthermore, by using delivery systems and mycotechnologies
17 disclosed herein instead of relying on serendipitous sporefalls, environmental
18 designers can greatly benefit by establishing, strengthening or steering the
19 course of habitat evolution in a fashion that is both environmentally sound
20 and/or economically profitable. In installing new parks, landscapes, forests,
21 arboretums, habitat oases and oasis-islands, space colonies, terrestrial

1 environments on this planet and on others, the insertion of purposely
2 designed 'fungal footprints' can dramatically improve the biodynamics of any
3 ecosystem.

4 It should be understood the foregoing detailed description is for
5 purposes of illustration rather than limitation of the scope of protection
6 accorded this invention, and therefore the description should be considered
7 illustrative, not exhaustive. The scope of protection is to be measured as
8 broadly as the invention permits. While the invention has been described in
9 connection with preferred embodiments, it will be understood that there is no
10 intention to limit the invention to those embodiments. On the contrary, it
11 will be appreciated that those skilled in the art, upon attaining an
12 understanding of the invention, may readily conceive of alterations to,
13 modifications of, and equivalents to the preferred embodiments without
14 departing from the principles of the invention, and it is intended to cover all
15 these alternatives, modifications and equivalents. Accordingly, the scope of
16 the present invention should be assessed as that of the appended claims and
17 any equivalents falling within the true spirit and scope of the invention.

18